

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 September 2006 (08.09.2006)

PCT

(10) International Publication Number
WO 2006/092143 A1

(51) International Patent Classification:
C07D 413/04 (2006.01) A61P 25/00 (2006.01)
A61K 31/5377 (2006.01)

(74) Agent: H. LUNDBECK A/S; Corporate Patents, Oetiliavej 9, DK-2500 Valby-Copenhagen (DK).

(21) International Application Number:
PCT/DK2006/000123

(22) International Filing Date: 2 March 2006 (02.03.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/658,428 3 March 2005 (03.03.2005) US
PA 200500321 3 March 2005 (03.03.2005) DK

(71) Applicant (for all designated States except US): H. LUNDBECK A/S [DK/DK]; Oetiliavej 9, DK-2500 Valby-Copenhagen (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): TORNØE, Christian, Wenzel [DK/DK]; Sverrigsgade 20, 2. Th., DK-2300 Copenhagen S (DK). KHANZHIN, Nikolay [RU/DK]; Teglgårdsvej 545, 2. Tv, DK-3050 Humlebæk (DK). RÖTTLÄNDER, Mario [DE/DK]; Svanemosen 10, DK-2670 Greve (DK). WATSON, William, Patrick [GB/DK]; Jyllingevej 81 St., DK-2720 Vanløse (DK). GREVE, Daniel, Rodriguez [DK/DK]; Kornvænget 20, DK-3660 Stenløse (DK).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED PYRIDINE DERIVATIVES



(57) Abstract: The present invention relates to pyridine derivatives of the general formula (I) and their use as openers of the KCNQ family potassium ion channels for the treatment of CNS disorders.

WO 2006/092143 A1

SUBSTITUTED PYRIDINE DERIVATIVES

FIELD OF THE INVENTION

The present invention relates to compounds, which are openers of the KCNQ family potassium ion channels. The compounds are useful in the treatment of disorders and diseases being responsive to opening of the KCNQ family potassium ion channels, one such disease is epilepsy.

10 BACKGROUND OF THE INVENTION

Ion channels are cellular proteins that regulate the flow of ions, including potassium, calcium, chloride and sodium into and out of cells. Such channels are present in all animal and human cells and affect a variety of processes including neuronal transmission, muscle contraction, and cellular secretion.

15

Humans have over 70 genes encoding potassium channel subtypes (Jentsch *Nature Reviews Neuroscience* 2000, 1, 21-30) with a great diversity with regard to both structure and function. Neuronal potassium channels, which are found in the brain, are primarily responsible for maintaining a negative resting membrane potential, as well as controlling membrane repolarisation following an action potential.

20

One subset of potassium channel genes is the KCNQ family. Mutations in four out of five KCNQ genes have been shown to underlie diseases including cardiac arrhythmias, deafness and epilepsy (Jentsch *Nature Reviews Neuroscience* 2000, 1, 21-30).

25

The KCNQ4 gene is thought to encode the molecular correlate of a potassium channel found in outer hair cells of the cochlea and in Type I hair cells of the vestibular apparatus, in which, mutations can lead to a form of inherited deafness.

30

KCNQ1 (KvLQT1) is co-assembled with the product of the KCNE1 (minimal K(+)-channel protein) gene in the heart to form a cardiac-delayed rectifier-like K(+) current. Mutations in this channel can cause one form of inherited long QT syndrome type 1 (LQT1), as well as being associated with a form of deafness (Robbins *Pharmacol Ther* 2001, 90, 1-19).

The genes KCNQ2 and KCNQ3 were discovered in 1988 and appear to be mutated in an inherited form of epilepsy known as benign familial neonatal convulsions (Rogawski *Trends in Neurosciences* 2000, 23, 393-398). The proteins encoded by the KCNQ2 and
5 KCNQ3 proteins are localised in the pyramidal neurons of the human cortex and hippocampus, regions of the brain associated with seizure generation and propagation (Cooper et al. *Proceedings National Academy of Science U S A* 2000, 97, 4914-4919).

KCNQ2 and KCNQ3 are two potassium channel subunits that form "M-currents" when
10 expressed in vitro. The M-current is a non-inactivating potassium current found in many neuronal cell types. In each cell type, it is dominant in controlling membrane excitability by being the only sustained current in the range of action potential initiation (Marrion *Annual Review Physiology* 1997, 59, 483-504). Modulation of the M-current has dramatic effects on neuronal excitability, for example activation of the current will reduce neuronal excitability.
15 Openers of these KCNQ channels, or activators of the M-current, will reduce excessive neuronal activity and may thus be of use in the treatment of seizures and other diseases and disorders characterised by excessive neuronal activity, such as neuronal hyperexcitability including convulsive disorders, epilepsy and neuropathic pain.

20 Retigabine (D-23129; *N*-(2-amino-4-(4-fluorobenzylamino)-phenyl) carbamic acid ethyl ester) and analogues thereof are disclosed in EP554543. Retigabine is an anti-convulsive compound with a broad spectrum and potent anticonvulsant properties, both *in vitro* and *in vivo*. It is active after oral and intraperitoneal administration in rats and mice in a range of anticonvulsant tests including: electrically induced seizures, seizures induced chemically by
25 pentylenetetrazole, picrotoxin and *N*-methyl-D-aspartate (NMDA) and in a genetic animal model, the DBA/2 mouse (Rostock et al. *Epilepsy Research* 1996, 23, 211-223). In addition, retigabine is active in the amygdala kindling model of complex partial seizures, further indicating that this compound has potential for anti-convulsive therapy. In clinical trials, retigabine has recently shown effectiveness in reducing the incidence of seizures in epileptic
30 patients (Bialer et al. *Epilepsy Research* 2002, 51, 31-71).

Retigabine has been shown to activate a K(+) current in neuronal cells and the pharmacology of this induced current displays concordance with the published

pharmacology of the M-channel, which recently was correlated to the KCNQ2/3 K(+) channel heteromultimer. This suggests that activation of KCNQ2/3 channels may be responsible for some of the anticonvulsant activity of this agent (Wickenden et al. *Molecular Pharmacology* 2000, 58, 591-600) – and that other agents working by the same
5 mechanism may have similar uses.

KCNQ 2 and 3 channels have also been reported to be upregulated in models of neuropathic pain (Wickenden et al. *Society for Neuroscience Abstracts* 2002, 454.7), and potassium channel modulators have been hypothesised to be active in both neuropathic pain and
10 epilepsy (Schroder et al. *Neuropharmacology* 2001, 40, 888-898).

Retigabine has also been shown to be beneficial in animal models of neuropathic pain (Blackburn-Munro and Jensen *European Journal of Pharmacology* 2003, 460, 109-116), and it is thus suggested that openers of KCNQ channels will be of use in treating pain
15 disorders including neuropathic pain.

The localisation of KCNQ channel mRNA is reported in brain and other central nervous system areas associated with pain (Goldstein et al. *Society for Neuroscience Abstracts* 2003, 53.8).
20

In addition to a role in neuropathic pain, the expression of mRNA for KCNQ 2-5 in the trigeminal and dorsal root ganglia and in the trigeminal nucleus caudalis implies that openers of these channels may also affect the sensory processing of migraine pain (Goldstein et al. *Society for Neuroscience Abstracts* 2003, 53.8).
25

Recent reports demonstrate that mRNA for KCNQ 3 and 5, in addition to that for KCNQ2, are expressed in astrocytes and glial cells. Thus KCNQ 2, 3 and 5 channels may help modulate synaptic activity in the CNS and contribute to the neuroprotective effects of KCNQ channel openers (Noda et al., *Society for Neuroscience Abstracts* 2003, 53.9).
30

Retigabine and other KCNQ modulators may thus exhibit protection against the neurodegenerative aspects of epilepsy, as retigabine has been shown to prevent limbic neurodegeneration and the expression of markers of apoptosis following kainic acid-induced

status epilepticus in the rat (Ebert et al. *Epilepsia* 2002, 43 Suppl 5, 86-95). This may have relevance for preventing the progression of epilepsy in patients, i.e. be anti-epileptogenic. Retigabine has also been shown to delay the progression of hippocampal kindling in the rat, a further model of epilepsy development (Tober et al. *European Journal Of Pharmacology* 5 1996, 303, 163-169).

It is thus suggested that these properties of retigabine and other KCNQ modulators may prevent neuronal damage induced by excessive neuronal activation, and such compounds may be of use in the treatment of neurodegenerative diseases, and be disease modifying (or 10 antiepileptogenic) in patients with epilepsy.

Given that anticonvulsant compounds such as benzodiazepines and chlormethiazole are used clinically in the treatment of the ethanol withdrawal syndrome and that other anticonvulsant compounds e.g. gabapentin, are very effective in animal models of this 15 syndrome (Watson et al. *Neuropharmacology* 1997, 36, 1369-1375), other anticonvulsant compounds such as KCNQ openers are thus expected to be effective in this condition.

mRNA for KCNQ 2 and 3 subunits are found in brain regions associated with anxiety and emotional behaviours such as bipolar disorder e.g. hippocampus and amygdala (Saganich et 20 al. *Journal of Neuroscience* 2001, 21, 4609-4624), and retigabine is reportedly active in some animal models of anxiety-like behaviour (Hartz et al. *Journal of Psychopharmacology* 2003, 17 suppl 3, A28,B16), and other clinically used anticonvulsant compounds are used in the treatment of bipolar disorder. Thus, KCNQ openers may be useful for the treatment of anxiety disorders and bipolar disorder.

25

WO 200196540 discloses the use of modulators of the M-current formed by expression of KCNQ2 and KCNQ3 genes for insomnia, while WO 2001092526 discloses that modulators of KCNQ5 can be utilized for the treatment of sleep disorders.

30 WO01/022953 describes the use of retigabine for prophylaxis and treatment of neuropathic pain such as allodynia, hyperalgesic pain, phantom pain, neuropathic pain related to diabetic neuropathy and neuropathic pain related to migraine.

WO02/049628 describes the use of retigabine for the treatment of anxiety disorders such as anxiety, generalized anxiety disorder, panic anxiety, obsessive compulsive disorder, social phobia, performance anxiety, post-traumatic stress disorder, acute stress reaction, adjustment disorders, hypochondriacal disorders, separation anxiety disorder, agoraphobia
5 and specific phobias.

- WO97/15300 describes the use of retigabine for the treatment of neurodegenerative disorders such as Alzheimer's disease; Huntington's chorea; sclerosis such as multiple sclerosis and amyotrophic lateral sclerosis; Creutzfeld-Jakob disease; Parkinson's disease;
10 encephalopathies induced by AIDS or infection by rubella viruses, herpes viruses, borrelia and unknown pathogens; trauma-induced neurodegenerations; neuronal hyperexcitation states such as in medicament withdrawal or intoxication; and neurodegenerative diseases of the peripheral nervous system such as polyneuropathies and polyneuritides.
- 15 KCNQ channel openers have also been found to be effective in the treatment of stroke, therefore it can be expected that selective KCNQ openers are effective in the treatment of stroke (Schroder et al., Pflugers Arch., 2003; 446(5): 607-16; Cooper and Jan, Arch Neurol., 2003, 60(4):496-500; Jensen, CNS Drug Rev., 2002, 8(4):353-60).
- 20 KCNQ channels have been shown to be expressed in dopaminergic and cholinergic circuits in the brain that are associated with the brain's reward system, particularly the ventral tegmental area (Cooper et al., J Neurosci, 2001, 21, 9529-9540). Therefore, KCNQ channel openers are expected to be effective in hyperexcitability disorders that involve the brain's reward system such as cocaine abuse, nicotine withdrawal and ethanol withdrawal.
- 25 Potassium channels comprised of the KCNQ4 subunits are expressed in the inner ear (Kubisch et al., Cell., 1999 Feb 5;96(3):437-46) and opening of these channels is therefore expected to treat tinnitus.
- 30 Hence, there is a great desire for novel compounds which are potent openers of the KCNQ family of potassium channels.

Also desired are novel compounds with improved properties relative to known compounds, which are openers of the KCNQ family potassium channels, such as retigabine.


Improvement of one or more of the following parameters is desired:

- 5 half-life, clearance, selectivity, interactions with other medications, bioavailability, potency, formulability, chemical stability, metabolic stability, membrane permeability, solubility and therapeutic index. The improvement of such parameters may lead to improvements such as:
- an improved dosing regime by reducing the number of required doses a day,
 - ease of administration to patients on multiple medications,
 - reduced side effects,
 - 10 • enlarged therapeutic index,
 - improved tolerability or
 - improved compliance.

15 SUMMARY OF THE INVENTION

One object of the invention is the provision of compounds, which are potent openers of the KCNQ family potassium channels.

- The compounds of the invention are substituted pyridine derivatives of the below formula I
20 as the free base or a salt thereof



wherein

R¹, R², R³ and q are as defined below.

25

The invention provides a compound of formula I for use as a medicament.

The invention provides a pharmaceutical composition comprising a compound of formula I and a pharmaceutically acceptable carrier or diluent.

The invention provides the use of a compound of formula I for the preparation of a
5 medicament for the treatment of seizure disorders, anxiety disorders, neuropathic pain and migraine pain disorders or neurodegenerative disorders.

The invention furthermore concerns the use of a compound of formula I in a method of treatment of seizure disorders, anxiety disorders, neuropathic pain and migraine pain
10 disorders or neurodegenerative disorders.

DEFINITION OF SUBSTITUENTS

The term "heteroatom" refers to a nitrogen, oxygen or sulphur atom.

15

"Halogen" means fluoro, chloro, bromo or iodo. "Halo" means halogen.

"Cyano" designates



20 which is attached to the remainder of the molecule via the carbon atom.

The expression "C₁₋₆-alk(en/yn)yl" means C₁₋₆-alkyl, C₂₋₆-alkenyl or C₂₋₆-alkynyl.

The term "C₁₋₆-alkyl" refers to a branched or unbranched alkyl group having from one to six carbon atoms, including but not limited to methyl, ethyl, prop-1-yl, prop-2-yl, 2-methyl-
25 prop-1-yl, 2-methyl-prop-2-yl, 2,2-dimethyl-prop-1-yl, but-1-yl, but-2-yl, 3-methyl-but-1-yl, 3-methyl-but-2-yl, pent-1-yl, pent-2-yl, pent-3-yl, hex-1-yl, hex-2-yl and hex-3-yl.

The term "C₂₋₆-alkenyl" refers to a branched or unbranched alkenyl group having from two to six carbon atoms and one double bond, including but not limited to ethenyl, propenyl, and butenyl.

30 The term "C₂₋₆-alkynyl" refers to a branched or unbranched alkynyl group having from two to six carbon atoms and one triple bond, including but not limited to ethynyl, propynyl and butynyl.

The expression "C₁₋₈-alk(en/yn)yl" means C₁₋₈-alkyl, C₂₋₈-alkenyl or C₂₋₈-alkynyl.

The term "C₁₋₈-alkyl" refers to a branched or unbranched alkyl group having from one to eight carbon atoms, including but not limited to methyl, ethyl, prop-1-yl, prop-2-yl, 2-methyl-prop-1-yl, 2-methyl-prop-2-yl, 2,2-dimethyl-prop-1-yl, but-1-yl, but-2-yl, 3-methyl-but-1-yl, 3-methyl-but-2-yl, pent-1-yl, pent-2-yl, pent-3-yl, hex-1-yl, hex-2-yl, hex-3-yl, 2-methyl-4,4-dimethyl-pent-1-yl and hept-1-yl.

The term "C₂₋₈-alkenyl" refers to a branched or unbranched alkenyl group having from two to eight carbon atoms and one double bond, including but not limited to ethenyl, propenyl, and butenyl.

- 10 The term "C₂₋₈-alkynyl" refers to a branched or unbranched alkynyl group having from two to eight carbon atoms and one triple bond, including but not limited to ethynyl, propynyl and butynyl.

The expression "C₃₋₈-cycloalk(en)yl" means C₃₋₈-cycloalkyl or C₃₋₈-cycloalkenyl.

- 15 The term "C₃₋₈-cycloalkyl" designates a monocyclic or bicyclic carbocycle having three to eight carbon atoms, including but not limited to cyclopropyl, cyclopentyl, cyclohexyl, bicycloheptyl such as 2-bicyclo[2.2.1]heptyl.

The term "C₃₋₈-cycloalkenyl" designates a monocyclic or bicyclic carbocycle having three to eight carbon atoms and one double bond, including but not limited to cyclopentenyl and

- 20 cyclohexenyl.

The term "C₃₋₈-heterocycloalk(en)yl" means C₃₋₈-heterocycloalkyl or C₃₋₈-heterocycloalkenyl.

- 25 The term "C₃₋₈-heterocycloalkyl" designates a monocyclic or bicyclic ring system wherein the ring is formed by 3 to 8 atoms selected from 2-7 carbon atoms and 1 or 2 heteroatoms independently selected from nitrogen, oxygen and sulphur atoms. Examples of C₃₋₈-heterocycloalkyls are pyrrolidine, azepan, morpholine, piperidine, piperazine and tetrahydrofuran.

- 30 The term "C₃₋₈-heterocycloalkenyl" designates a monocyclic or bicyclic ring system with one double bond, wherein the ring is formed by 3 to 8 atoms selected from 2-7 carbon atoms and 1 or 2 heteroatoms independently selected from nitrogen, oxygen and sulphur atoms. Examples of C₃₋₈-heterocycloalkenyls are dihydropyrrole, dihydrofuran and dihydrothiophene.

When C₃₋₈-heterocycloalk(en)yl comprises nitrogen then C₃₋₈-heterocycloalk(en)yl is attached to the remainder of the molecule via a carbon atom or nitrogen atom of the heterocyclic ring.

When C₃₋₈-heterocycloalk(en)yl does not comprise nitrogen then C₃₋₈-heterocycloalk(en)yl
5 is attached to the remainder of the molecule via a carbon atom of the heterocyclic ring.

The term "halo-C₁₋₆-alk(en/yn)yl" designates C₁₋₆-alk(en/yn)yl being substituted with halogen, including but not limited to trifluoromethyl.

Similarly, "halo-C₃₋₈-cycloalk(en)yl" designates C₃₋₈-cycloalk(en)yl being substituted with
10 halogen, including but not limited to chlorocyclopropane and chlorocyclohexane.
Similarly, "halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl" designates halo-C₃₋₈-cycloalk(en)yl being attached to the remainder of the molecule via C₁₋₆-alk(en/yn)yl.

The term "C₁₋₆-alk(en/yn)oxy" designates C₁₋₆-alk(en/yn)yl being attached to the
15 remainder of the molecule via an oxygen atom.

Similarly, "C₃₋₈-cycloalk(en)oxy" designates C₃₋₈-cycloalk(en)yl being attached to the remainder of the molecule via an oxygen atom.

In the expressions "C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl", "Aryl-C₁₋₆-alk(en/yn)yl", "Aryl-
20 C₃₋₈-cycloalk(en)yl", "Aryl-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl",
"C₃₋₈-heterocycloalk(en)yl-C₁₋₆-alk(en/yn)yl", "C₁₋₆-alk(en/yn)yl-C₃₋₈-heterocycloalk(en)yl-
C₁₋₆-alk(en/yn)yl", "Heteroaryl-C₁₋₆-alk(en/yn)yl", "Heteroaryl-C₃₋₈-cycloalk(en)yl",
"Heteroaryl-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl", "NR⁴R⁵-C₁₋₆-alk(en/yn)yl", "NR⁴R⁵-
C₃₋₈-cycloalk(en)yl", "NR⁴R⁵-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl", "C₃₋₈-cycloalk(en)yl-
25 C₁₋₆-alk(en/yn)oxy", "C₁₋₆-alk(en/yn)oxy-C₁₋₆-alk(en/yn)yl", "C₃₋₈-cycloalk(en)oxy-
C₁₋₆-alk(en/yn)yl" and "C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)oxy-C₁₋₆-alk(en/yn)yl" the
terms "C₁₋₆-alk(en/yn)yl", "C₃₋₈-cycloalk(en)yl", "Aryl", "C₃₋₈-heterocycloalk(en)yl",
"Heteroaryl", "C₁₋₆-alk(en/yn)oxy" and "C₃₋₈-cycloalk(en)oxy" are as defined above.

30 The term "Heteroaryl" refers to monocyclic or bicyclic heteroaromatic systems being selected from the group consisting of pyridine, thiophene, furan, pyrrole, pyrazole, triazole, tetrazole, oxazole, imidazole, thiazole, benzofuran, benzothiophene and indole.

The term Aryl designates monocyclic or bicyclic aromatic systems being selected from the group consisting of phenyl and naphthyl.

The term "optionally substituted Aryl-C₁₋₆-alk(en/yn)yl" designates Aryl-C₁₋₆-alk(en/yn)yl

5 wherein the Aryl moiety is optionally substituted, such as with 1, 2 or 3 substituents independently selected from the group consisting of halogen, cyano, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl, halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yloxy, C₃₋₈-cycloalk(en)yloxy and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yloxy.

10 Similarly, "optionally substituted Aryl-C₃₋₈-cycloalk(en)yl" designates Aryl-C₃₋₈-cycloalk(en)yl wherein the Aryl moiety is optionally substituted, such as with 1, 2 or 3 substituents independently selected from the group consisting of halogen, cyano,

C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl, halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl,


15 C₁₋₆-alk(en/yn)yloxy, C₃₋₈-cycloalk(en)yloxy and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yloxy.

Similarly, "optionally substituted Aryl-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl" designates Aryl-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl wherein the Aryl moiety is optionally substituted, such as with 1, 2 or 3 substituents independently selected from the group consisting of halogen, cyano, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl, halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yloxy, C₃₋₈-cycloalk(en)yloxy and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yloxy.

DESCRIPTION OF THE INVENTION

25 The present invention relates to substituted pyridine derivatives which are openers of KCNQ potassium channels.

The present invention relates to a compound represented by the general formula I:



wherein

q is 0 or 1;

each of R¹ and R² is independently selected from the group consisting of halogen, cyano,

- 5 C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl, halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yoxy, C₃₋₈-cycloalk(en)yoxy and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yoxy; and

R³ is selected from the group consisting of C₁₋₈-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl,

- 10 C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, optionally substituted Aryl-C₁₋₆-alk(en/yn)yl, optionally substituted Aryl-C₃₋₈-cycloalk(en)yl, optionally substituted Aryl-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yl-C₃₋₈-heterocycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₃₋₈-heterocycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yl-C₃₋₈-heterocycloalk(en)yl-C₁₋₆-alk(en/yn)yl, Heteroaryl-C₁₋₆-alk(en/yn)yl, Heteroaryl-C₃₋₈-cycloalk(en)yl, Heteroaryl-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, NR⁴R⁵-C₁₋₆-alk(en/yn)yl, NR⁴R⁵-C₃₋₈-cycloalk(en)yl, NR⁴R⁵-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yoxy-C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yoxy-C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yoxy-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl and halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl;

- 20 wherein

each of R⁴ and R⁵ is independently selected from the group consisting of hydrogen, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl; as the free base or salts thereof.

- 25 In one embodiment of the compound of formula I, q is 0;
in another embodiment of the compound of formula I, q is 1.

- In a further embodiment of the compound of formula I each of R^1 and R^2 is independently selected from the group consisting of halogen, cyano, halo- C_{1-6} -alk(en/yn)yl, halo- C_{3-8} -cycloalk(en)yl, halo- C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl, C_{1-6} -alk(en/yn)yoxy, C_{3-8} -cycloalk(en)yoxy and C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yoxy;
- 5 in another embodiment each of R^1 and R^2 is independently selected from the group consisting of halogen, cyano, C_{1-6} -alk(en/yn)yl, C_{3-8} -cycloalk(en)yl, C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl, C_{1-6} -alk(en/yn)yoxy, C_{3-8} -cycloalk(en)yoxy and C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yoxy;
- 10 in another embodiment each of R^1 and R^2 is independently selected from the group consisting of halogen, cyano and C_{1-6} -alk(en/yn)yl and C_{1-6} -alk(en/yn)yoxy;
- in another embodiment each of R^1 and R^2 is independently selected from the group consisting of C_{1-6} -alk(en/yn)yl, C_{3-8} -cycloalk(en)yl and C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl;
- in another embodiment R^1 is C_{1-6} -alk(en/yn)yl, such as methyl;
- 15 in another embodiment R^2 is C_{1-6} -alk(en/yn)yl, such as methyl;
- in another embodiment R^1 is C_{1-6} -alk(en/yn)yoxy, such as methoxy and R^2 is halogen;
- in another embodiment R^1 is halogen and R^2 is C_{1-6} -alk(en/yn)yoxy, such as methoxy.
- Typically, both R^1 and R^2 are C_{1-6} -alk(en/yn)yl, such as methyl.
- 20 In a further embodiment of the compound of formula I, R^3 is selected from the group consisting of C_{1-6} -alk(en/yn)yl- C_{3-8} -heterocycloalk(en)yl- C_{1-6} -alk(en/yn)yl, C_{3-8} -heterocycloalk(en)yl- C_{1-6} -alk(en/yn)yl, C_{1-6} -alk(en/yn)yl- C_{3-8} -heterocycloalk(en)yl- C_{1-6} -alk(en/yn)yl, NR^4R^5 - C_{1-6} -alk(en/yn)yl, NR^4R^5 - C_{3-8} -cycloalk(en)yl, NR^4R^5 - C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl, C_{1-6} -alk(en/yn)yoxy- C_{1-6} -alk(en/yn)yl,
- 25 C_{3-8} -cycloalk(en)yoxy- C_{1-6} -alk(en/yn)yl, C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yoxy- C_{1-6} -alk(en/yn)yl, halo- C_{1-6} -alk(en/yn)yl, halo- C_{3-8} -cycloalk(en)yl and halo- C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl;
- in another embodiment R^3 is selected from the group consisting of C_{1-8} -alk(en/yn)yl, C_{3-8} -cycloalk(en)yl, C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl, optionally substituted Aryl- C_{1-6} -alk(en/yn)yl, optionally substituted Aryl- C_{3-8} -cycloalk(en)yl, optionally substituted Aryl- C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl, Heteroaryl- C_{1-6} -alk(en/yn)yl, Heteroaryl- C_{3-8} -cycloalk(en)yl, Heteroaryl- C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl, NR^4R^5 .
- 30

C₁₋₆-alk(en/yn)yl, NR⁴R⁵-C₃₋₈-cycloalk(en)yl and NR⁴R⁵-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl;

in another embodiment R³ is selected from the group consisting of C₁₋₈-alk(en/yn)yl,

C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, optionally substituted Aryl-C₁₋₆-alk(en/yn)yl,

- 5 optionally substituted Aryl-C₃₋₈-cycloalk(en)yl, Heteroaryl-C₁₋₆-alk(en/yn)yl and NR⁴R⁵-C₁₋₆-alk(en/yn)yl;

in another embodiment R³ is selected from the group consisting of C₁₋₈-alk(en/yn)yl,

C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, optionally substituted Aryl-

C₁₋₆-alk(en/yn)yl, optionally substituted Aryl-C₃₋₈-cycloalk(en)yl, optionally substituted

- 10 Aryl-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, Heteroaryl-C₁₋₆-alk(en/yn)yl, Heteroaryl-C₃₋₈-cycloalk(en)yl and Heteroaryl-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl.

Typically, R³ is selected from the group consisting of C₁₋₈-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, optionally substituted Aryl-C₁₋₆-alk(en/yn)yl, optionally substituted Aryl-C₃₋₈-cycloalk(en)yl and Heteroaryl-C₁₋₆-alk(en/yn)yl.

- 15 To further illustrate without limiting the invention, an embodiment of R³ is C₁₋₈-alk(en/yn)yl;

another embodiment of R³ is C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl;

another embodiment of R³ is optionally substituted Aryl-C₁₋₆-alk(en/yn)yl;

another embodiment of R³ is optionally substituted Aryl-C₃₋₈-cycloalk(en)yl;

- 20 another embodiment of R³ is Heteroaryl-C₁₋₆-alk(en/yn)yl.

In a further embodiment of the compound of formula I, each of R⁴ and R⁵ is independently selected from the group consisting of C₃₋₈-cycloalk(en)yl and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl;

- 25 in another embodiment each of R⁴ and R⁵ is independently selected from the group consisting of C₁₋₆-alk(en/yn)yl and hydrogen;

in another embodiment both R⁴ and R⁵ are C₁₋₆-alk(en/yn)yl;

in another embodiment both R⁴ and R⁵ are hydrogen.

- 30 In a further embodiment of the compound of formula I, any Heteroaryl, which is mentioned either alone or as a part of a larger substituent is selected from the group consisting of pyridine, furan, pyrrole, pyrazole, triazole, tetrazole, oxazole, imidazole, thiazole,

benzofuran, benzothiophene and indole; in another embodiment any Heteroaryl, which is mentioned either alone or as a part of a larger substituent is thiophene.

In a further embodiment of the compound of formula I, any Aryl, which is mentioned either
5 alone or as a part of a larger substituent is phenyl;
in another embodiment any Aryl, which is mentioned either alone or as a part of a larger substituent is naphthyl.

In a further embodiment of the compound of formula I, any optionally substituted Aryl,
10 which is mentioned either alone or as a part of a larger substituent, may be substituted with 1 or 2 substituents.

To further illustrate without limiting the invention an embodiment concerns such compounds of formula I, wherein any optionally substituted Aryl which is mentioned either alone or as a part of a larger substituent is not substituted;
15 in another embodiment any optionally substituted Aryl which is mentioned either alone or as a part of a larger substituent is substituted with 1 substituent;
in another embodiment any optionally substituted Aryl which is mentioned either alone or as a part of a larger substituent is substituted with 2 substituents.

20 In a further embodiment of the compound of formula I, any optionally substituted Aryl, which is mentioned either alone or as a part of a larger substituent, may be substituted with substituents selected from the group consisting of cyano, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl, halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yloxy and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yloxy;
25 in another embodiment any optionally substituted Aryl which is mentioned either alone or as a part of a larger substituent may be substituted with substituents selected from the group consisting of halogen, C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl and C₁₋₆-alk(en/yn)yloxy.
To further illustrate without limiting the invention an embodiment concerns such compounds of formula I, wherein any optionally substituted Aryl which is mentioned either
30 alone or as a part of a larger substituent is substituted with halogen;
in another embodiment any optionally substituted Aryl which is mentioned either alone or as a part of a larger substituent is substituted with C₁₋₆-alk(en/yn)yl;

in another embodiment any optionally substituted Aryl which is mentioned either alone or as a part of a larger substituent is substituted with halo-C₁₋₆-alk(en/yn)yl;

in another embodiment any optionally substituted Aryl which is mentioned either alone or as a part of a larger substituent is substituted with C₁₋₆-alk(en/yn)yl-oxy.

5

A further embodiment concerns a compound of formula I as the free base or a salt thereof, said compound is selected from the compounds of the following scheme:

Example No.	Compound name
1aa	(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-carbamic acid benzyl ester
1ab	(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-carbamic acid 2-chloro-benzyl ester
1ac	2-(4-Chloro-phenyl)-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide
1ad	2-Phenyl-cyclopropanecarboxylic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide
1ae	N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-thiophen-2-yl-acetamide
1af	3-Cyclohexyl-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-propionamide
1ag	(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-carbamic acid isobutyl ester
1ah	3-(3-Chloro-phenyl)-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-propionamide
1ai	N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(3,5-dimethyl-phenyl)-acetamide
1aj	N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-3-p-tolyl-propionamide
1ak	2-(3-Chloro-phenyl)-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide
1al	2-(3,4-Dichloro-phenyl)-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide
1am	N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-thiophen-3-yl-acetamide
1an	N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-p-tolyl-acetamide
1ao	2-(3-Bromo-phenyl)-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide
1ap	N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(3-trifluoromethyl-phenyl)-acetamide
1aq	N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-phenyl-acetamide
1ar	3,5,5-Trimethyl-hexanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide
1as	Octanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide
1at	N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-naphthalen-2-yl-acetamide
1au	Heptanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide
1av	N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(3,4-dimethyl-phenyl)-acetamide

law	2-(Cyclohex-1-enyl)- <i>N</i> -(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide
lax	<i>N</i> -(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(4-methoxy-3-methyl-phenyl)-acetamide
lay	<i>N</i> -(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(4-methoxy-phenyl)-acetamide
laz	<i>N</i> -(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-3-(4-methoxy-phenyl)-propionamide
lba	<i>N</i> -(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2- <i>m</i> -tolyl-acetamide
lbb	<i>N</i> -(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(4-fluoro-phenyl)-acetamide
lbc	<i>N</i> -(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-3,3-dimethyl-butyramide
lbd	<i>N</i> -(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(3-fluoro-phenyl)-acetamide
lbe	2-Bicyclo[2.2.1]hept-2-yl- <i>N</i> -(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide
lbf	2-(3,4-Difluoro-phenyl)- <i>N</i> -(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide
lbg	4-Methyl-pentanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide
lbh	2-(Cyclopent-2-enyl)- <i>N</i> -(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide
lbi	2-Cyclohexyl- <i>N</i> -(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide
lbj	5-Methyl-hexanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide
lbk	2-Cyclopentyl- <i>N</i> -(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide
lbl	3-Cyclopentyl- <i>N</i> -(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-propionamide
lbm	Hexanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide
lbn	<i>N</i> -(4-Chloro-2-methoxy-6-morpholin-4-yl-pyridin-3-yl)-2-cyclopentylacetamide
lbo	<i>N</i> -(2-Chloro-4-methoxy-6-morpholin-4-yl-pyridin-3-yl)-2-cyclopentylacetamide
lbp	<i>N</i> -(2-Chloro-4-methoxy-6-morpholin-4-yl-pyridin-3-yl)-3,3-dimethylbutyramide
lbq	<i>N</i> -(4-Chloro-2-methoxy-6-morpholin-4-yl-pyridin-3-yl)-3,3-dimethylbutyramide
lbr	<i>N</i> -(4-Chloro-2-methoxy-6-morpholin-4-yl-pyridin-3-yl)-propionamide

Each of these compounds is considered a specific embodiment and may be subjected to individual claims.

The present invention also comprises salts of the compounds of the invention, typically,
5 pharmaceutically acceptable salts. The salts of the invention include acid addition salts, metal salts, ammonium and alkylated ammonium salts.

The salts of the invention are preferably acid addition salts. The acid addition salts of the invention are preferably pharmaceutically acceptable salts of the compounds of the
10 invention formed with non-toxic acids. Acid addition salts include salts of inorganic acids as well as organic acids. Examples of suitable inorganic acids include hydrochloric,

hydrobromic, hydroiodic, phosphoric, sulfuric, sulfamic, nitric acids and the like. Examples of suitable organic acids include formic, acetic, trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, glycolic, itaconic, lactic, methanesulfonic, maleic, malic, malonic, mandelic, oxalic, picric, pyruvic, salicylic, succinic, methane sulfonic, ethanesulfonic, tartaric, ascorbic, pamoic, bismethylene salicylic, ethanedisulfonic, gluconic, citraconic, aspartic, stearic, palmitic, EDTA, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, p-toluenesulfonic acids, theophylline acetic acids, as well as the 8-halotheophyllines, for example 8-bromotheophylline and the like. Further examples of pharmaceutical acceptable inorganic or organic acid addition salts include the pharmaceutically acceptable salts listed in J. Pharm. Sci. 1977,66,2, which is incorporated herein by reference.

Also intended as acid addition salts are the hydrates, which the present compounds, are able to form.

Examples of metal salts include lithium, sodium, potassium, magnesium salts and the like.

Examples of ammonium and alkylated ammonium salts include ammonium, methyl-, dimethyl-, trimethyl-, ethyl-, hydroxyethyl-, diethyl-, *n*-butyl-, *sec*-butyl-, *tert*-butyl-, tetramethylammonium salts and the like.

Further, the compounds of this invention may exist in unsolvated as well as in solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of this invention.

The compounds of the present invention may have one or more asymmetric centre and it is intended that any optical isomers (i.e. enantiomers or diastereomers), as separated, pure or partially purified optical isomers and any mixtures thereof including racemic mixtures, i.e. a mixture of stereoisomers, are included within the scope of the invention.

Racemic forms can be resolved into the optical antipodes by known methods, for example, by separation of diastereomeric salts with an optically active acid, and liberating the

optically active amine compound by treatment with a base. Another method for resolving racemates into the optical antipodes is based upon chromatography on an optically active matrix. Racemic compounds of the present invention can also be resolved into their optical antipodes, e.g. by fractional crystallization. The compounds of the present invention may
5 also be resolved by the formation of diastereomeric derivatives. Additional methods for the resolution of optical isomers, known to those skilled in the art, may be used. Such methods include those discussed by J. Jaques, A. Collet and S. Wilen in "Enantiomers, Racemates, and Resolutions", John Wiley and Sons, New York (1981). Optically active compounds can also be prepared from optically active starting materials, or by stereoselective synthesis.

10

Furthermore, when a double bond or a fully or partially saturated ring system is present in the molecule, geometric isomers may be formed. It is intended that any geometric isomers, as separated, pure or partially purified geometric isomers or mixtures thereof are included within the scope of the invention. Likewise, molecules having a bond with restricted
15 rotation may form geometric isomers. These are also intended to be included within the scope of the present invention.

Furthermore, some of the compounds of the present invention may exist in different tautomeric forms and it is intended that any tautomeric forms that the compounds are able to
20 form are included within the scope of the present invention.

The invention also encompasses prodrugs of the present compounds, which on administration undergo chemical conversion by metabolic processes before becoming pharmacologically active substances. In general, such prodrugs will be functional
25 derivatives of the compounds of the general formula I, which are readily convertible in vivo into the required compound of the formula I. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

30 The invention also encompasses active metabolites of the present compounds.

The compounds according to the invention have affinity for the KCNQ2 receptor subtype with an EC_{50} of less than 15000nM such as less than 10000nM as measured by the test

"Relative efflux through the KCNQ2 channel" which is described below. One embodiment concerns such compounds of formula I having affinity for the KCNQ2 receptor subtype with an EC_{50} of less than 2000nM such as less than 1500nM as measured by the test

5 "Relative efflux through the KCNQ2 channel" which is described below. To further illustrate without limiting the invention an embodiment concerns such compounds having affinity for the KCNQ2 receptor subtype with an EC_{50} of less than 200nM such as less than 150nM as measured by the test "Relative efflux through the KCNQ2 channel" which is described below.

10 One embodiment concerns such compounds of formula I having an ED_{50} of less than 15 mg/kg in the test "Maximum electroshock" which is described below. To further illustrate without limiting the invention, an embodiment concerns such compounds having an ED_{50} of less than 5 mg/kg in the test "Maximum electroshock" which is described below.

15 One embodiment concerns such compounds of formula I having an ED_{50} of less than 5 mg/kg in the "Electrical seizure -threshold test" and "Chemical seizure -threshold test" which is described below.

20 One embodiment concerns such compounds of formula I having few or clinically insignificant side effects. Some of the compounds according to the invention are thus tested in models of the unwanted sedative, hypothermic and ataxic actions.

25 One embodiment concerns such compounds of formula I having a large therapeutic index between anticonvulsant efficacy and side-effects such as impairment of locomotor activity or ataxic effects as measured by performance on a rotating rod. Such compounds will expectedly be well tolerated in patients permitting high doses to be used before side effects are seen. Thereby compliance with the therapy will expectedly be good and administration of high doses may be permitted making the treatment more efficacious in patients who would otherwise have side effects with other medications.

30

As already mentioned, the compounds according to the invention have effect on potassium channels of the KCNQ family, in particular the KCNQ2 subunit, and they are thus considered useful for increasing ion flow in a voltage-dependent potassium channel in a

mammal such as a human. The compounds of the invention are considered applicable in the treatment of a disorder or disease being responsive to an increased ion flow in a potassium channel such as the KCNQ family potassium ion channels. Such disorder or disease is preferably a disorder or disease of the central nervous system.

5

In one aspect, the compounds of the invention may be administered as the only therapeutically effective compound.

In another aspect the compounds of the invention may be administered as a part of a combination therapy, i.e. the compounds of the invention may be administered in

10 combination with other therapeutically effective compounds having e.g. anti-convulsive properties. The effects of such other compounds having anti-convulsive properties may include but not be limited to activities on:

- ion channels such as sodium, potassium, or calcium channels
- the excitatory amino acid systems e.g. blockade or modulation of NMDA receptors
- 15 • the inhibitory neurotransmitter systems e.g. enhancement of GABA release, or blockade of GABA-uptake or
- membrane stabilisation effects.

Current anti-convulsive medications include, but are not limited to, tiagabine, carbamazepine, sodium valproate, lamotrigine, gabapentin, pregabalin, ethosuximide, 20 levetiracetam, phenytoin, topiramate, zonisamide as well as members of the benzodiazepine and barbiturate class.

An aspect of the invention provides a compound of formula I free base or a salt thereof for use as a medicament.

25

In one embodiment, the invention relates to the use of a compound of formula I free base or a salt thereof in a method of treatment.

An embodiment of the invention provides a pharmaceutical composition comprising a 30 compound of formula I free base or a salt thereof and a pharmaceutically acceptable carrier or diluent. The composition may comprise any of the embodiments of formula I as described above.

A further embodiment of the invention relates to the use of a compound of formula I free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of a disease or disorder wherein a KCNQ potassium channel opener such as a KCNQ2 potassium channel opener is beneficial. Typically, such disorder or disease is selected from the group consisting of seizure disorders, anxiety disorders, neuropathic pain and migraine pain disorders, neurodegenerative disorders, stroke, cocaine abuse, nicotine withdrawal, ethanol withdrawal and tinnitus.

A further embodiment of the invention relates to the use of a compound of formula I free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of seizure disorders.

Typically, the seizure disorders to be treated are selected from the group consisting of acute seizures, convulsions, status epilepticus and epilepsy such as epileptic syndromes and epileptic seizures.

A further embodiment of the invention relates to the use of a compound of formula I free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of anxiety disorders.

Typically, the anxiety disorders to be treated are selected from the group consisting of anxiety and disorders and diseases related to panic attack, agoraphobia, panic disorder with agoraphobia, panic disorder without agoraphobia, agoraphobia without history of panic disorder, specific phobia, social phobia and other specific phobias, obsessive-compulsive disorder, posttraumatic stress disorder, acute stress disorders, generalized anxiety disorder, anxiety disorder due to general medical condition, substance-induced anxiety disorder, separation anxiety disorder, adjustment disorders, performance anxiety, hypochondriacal disorders, anxiety disorder due to general medical condition and substance-induced anxiety disorder and anxiety disorder not otherwise specified.

A further embodiment of the invention relates to the use of a compound of formula I free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of neuropathic pain and migraine pain disorders.

Typically, the neuropathic pain and migraine pain disorders to be treated are selected from the group consisting of allodynia, hyperalgesic pain, phantom pain, neuropathic pain related

to diabetic neuropathy, neuropathic pain related to trigeminal neuralgia and neuropathic pain related to migraine.

5 A further embodiment of the invention relates to the use of a compound of formula I free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of neurodegenerative disorders.

Typically the neurodegenerative disorders to be treated are selected from the group consisting of Alzheimer's disease, Huntington's chorea, multiple sclerosis, amyotrophic lateral sclerosis, Creutzfeld-Jakob disease, Parkinson's disease, encephalopathies induced
10 by AIDS or infection by rubella viruses, herpes viruses, borrelia and unknown pathogens, trauma-induced neurodegenerations, neuronal hyperexcitation states such as in medicament withdrawal or intoxication and neurodegenerative diseases of the peripheral nervous system such as polyneuropathies and polyneuritides.

15 A further embodiment of the invention relates to the use of a compound of formula I free base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of bipolar disorders.

A further embodiment of the invention relates to the use of a compound of formula I free
20 base or a salt thereof for the preparation of a pharmaceutical composition for the treatment of sleep disorders; such as insomnia.

The term "treatment" as used herein in connection with a disease or disorders includes also prevention, inhibition and amelioration as the case may be.

25

The invention provides compounds showing effect in at least one of the following tests:

- "Relative efflux through the KCNQ2 channel"

Which is a measure of the potency of the compound at the target channel

- "Maximum electroshock"

30 Which is a measure of seizures induced by non-specific CNS stimulation by electrical means

- "Pilocarpine induced seizures"

Seizures induced by pilocarpine are often difficult to treat with many existing antiseizure medications and so reflect a model of "drug resistant seizures"

- "Electrical seizure-threshold tests" and "Chemical seizure-threshold tests"

These models measure the threshold at which seizures are initiated, thus being models that detect whether compounds could delay seizure initiation.

- "Amygdala kindling"

Which is used as a measure of disease progression, as in normal animals the seizures in this model get more severe as the animal receives further stimulations.

- "Electrophysiological patch-clamp recordings in CHO cells" and

"electrophysiological recordings of KCNQ2, KCNQ2/KCNQ3 or KCNQ5 channels in oocytes"

In these tests voltage-activated KCNQ2, KCNQ2/KCNQ3 or KCNQ5 currents are recorded.

PHARMACEUTICAL COMPOSITIONS

The present invention also relates to a pharmaceutical composition. The compounds of the invention as the free base or salts thereof may be administered alone or in combination with pharmaceutically acceptable carriers or diluents, in either single or multiple doses. The pharmaceutical compositions according to the invention may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 19 Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

The pharmaceutical compositions may be specifically formulated for administration by any suitable route such as the oral, rectal, nasal, pulmonary, topical (including buccal and sublingual), transdermal, intracisternal, intraperitoneal, vaginal and parenteral (including subcutaneous, intramuscular, intrathecal, intravenous and intradermal) route, the oral route being preferred. It will be appreciated that the preferred route will depend on the general condition and age of the subject to be treated, the nature of the disorder or disease to be treated and the active ingredient chosen.

The pharmaceutical compositions formed by combining the compound of the invention and the pharmaceutical acceptable carriers are then readily administered in a variety of dosage forms suitable for the disclosed routes of administration. The formulations may conveniently be presented in unit dosage form by methods known in the art of pharmacy.

5

The compounds of this invention are generally utilized as the free base or as a pharmaceutically acceptable salt thereof. One example is an acid addition salt of a compound having the utility of a free base. When a compound of the invention contains a free base such salts are prepared in a conventional manner by treating a solution or
10 suspension of a free base of the invention with a chemical equivalent of a pharmaceutically acceptable acid. Representative examples are mentioned above.

Pharmaceutical compositions for oral administration may be solid or liquid. Solid dosage forms for oral administration include e.g. capsules, tablets, dragees, pills, lozenges,
15 powders, granules and tablette e.g. placed in a hard gelatine capsule in powder or pellet form or e.g. in the form of a troche or lozenge. Where appropriate, pharmaceutical compositions for oral administration may be prepared with coatings such as enteric coatings or they can be formulated so as to provide controlled release of the active ingredient such as sustained or prolonged release according to methods well known in the art. Liquid dosage
20 forms for oral administration include e.g. solutions, emulsions, suspensions, syrups and elixirs.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules or tablets, each containing a predetermined amount of the
25 active ingredient, and which may include a suitable excipient. Furthermore, the orally available formulations may be in the form of a powder or granules, a solution or suspension in an aqueous or non-aqueous liquid, or an oil-in-water or water-in-oil liquid emulsion.

Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous
30 solution and various organic solvents. Examples of solid carriers are lactose, terra alba, sucrose, cyclodextrin, talc, gelatine, agar, pectin, acacia, magnesium stearate, stearic acid, lower alkyl ethers of cellulose, corn starch, potato starch, gums and the like. Examples of

25

liquid carriers are syrup, peanut oil, olive oil, phospho lipids, fatty acids, fatty acid amines, polyoxyethylene and water.

- The carrier or diluent may include any sustained release material known in the art, such as
- 5 glyceryl monostearate or glyceryl distearate, alone or mixed with a wax.

Any adjuvants or additives usually used for such purposes such as colourings, flavourings, preservatives etc. may be used provided that they are compatible with the active ingredients.

- 10 The amount of solid carrier may vary but will usually be from about 25 mg to about 1 g. If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatine capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.
- 15 Tablets may be prepared by mixing the active ingredient with ordinary adjuvants or diluents and subsequently compressing the mixture in a conventional tableting machine.

- Pharmaceutical compositions for parenteral administration include sterile aqueous and nonaqueous injectable solutions, dispersions, suspensions or emulsions as well as sterile
- 20 powders to be reconstituted in sterile injectable solutions or dispersions prior to use. Depot injectable formulations are also contemplated as being within the scope of the present invention.

- For parenteral administration, solutions of the compound of the invention in sterile aqueous
- 25 solution, aqueous propylene glycol, aqueous vitamin E or sesame or peanut oil may be employed. Such aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. The aqueous solutions are particularly suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. The sterile aqueous media employed are all readily available by standard
- 30 techniques known to those skilled in the art.

Solutions for injections may be prepared by dissolving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution

to the desired volume, sterilising the solution and filling it in suitable ampoules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

- 5 Other suitable administration forms include suppositories, sprays, ointments, cremes, gels, inhalants, dermal patches, implants, etc.

A typical oral dosage is in the range of from about 0.001 to about 100 mg/kg body weight per day, preferably from about 0.01 to about 50 mg/kg body weight per day, and more
10 preferred from about 0.05 to about 10 mg/kg body weight per day administered in one or more dosages such as 1 to 3 dosages. The exact dosage will depend upon the frequency and mode of administration, the sex, age, weight and general condition of the subject treated, the nature and severity of the disorder or disease treated and any concomitant diseases to be treated and other factors evident to those skilled in the art.

15

The formulations may conveniently be presented in unit dosage form by methods known to those skilled in the art. A typical unit dosage form for oral administration one or more times per day such as 1 to 3 times per day may contain from 0.01 to about 1000 mg, such as about 0.01 to 100 mg, preferably from about 0.05 to about 500 mg, and more preferred from about
20 0.5 mg to about 200 mg.

For parenteral routes such as intravenous, intrathecal, intramuscular and similar administration, typically doses are in the order of about half the dose employed for oral administration.

25

Typical examples of recipes for the formulation of the invention are as follows:

- 1) Tablets containing 5.0 mg of a compound of the invention calculated as the free base:
- | | | |
|----|---------------------------|--------|
| 30 | Compound of the invention | 5.0 mg |
| | Lactose | 60 mg |
| | Maize starch | 30 mg |
| | Hydroxypropylcellulose | 2.4 mg |

Microcrystalline cellulose	19.2 mg
Croscarmellose Sodium Type A	2.4 mg
Magnesium stearate	0.84 mg

- 5 2) Tablets containing 0.5 mg of a compound of the invention calculated as the free base:

Compound of the invention	0.5 mg
Lactose	46.9 mg
Maize starch	23.5 mg
10 Povidone	1.8 mg
Microcrystalline cellulose	14.4 mg
Croscarmellose Sodium Type A	1.8 mg
Magnesium stearate	0.63 mg

- 15 3) Syrup containing per millilitre:

Compound of the invention	25 mg
Sorbitol	500 mg
Hydroxypropylcellulose	15 mg
Glycerol	50 mg
20 Methyl-paraben	1 mg
Propyl-paraben	0.1 mg
Ethanol	0.005 mL
Flavour	0.05 mg
Saccharin sodium	0.5 mg
25 Water	ad 1 mL

- 4) Solution for injection containing per millilitre:

Compound of the invention	0.5 mg
Sorbitol	5.1 mg
30 Acetic Acid	0.05 mg
Saccharin sodium	0.5 mg
Water	ad 1 mL


By the expression a compound of the invention is meant any one of the embodiments of formula I as described herein.

5 In a further aspect the present invention relates to a method of preparing a compound of the invention as described in the following.

Preparation of the compounds of the invention

10

The present invention relates to a compound represented by the general formula I:



wherein

15 q is 0 or 1;

each of R¹ and R² is independently selected from the group consisting of halogen, cyano, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl, halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl,

20 C₁₋₆-alk(en/yn)yloxy, C₃₋₈-cycloalk(en)yloxy and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yloxy;
and

25 R^3 is selected from the group consisting of C_{1-8} -alk(en/yn)yl, C_{3-8} -cycloalk(en)yl, C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl, optionally substituted Aryl- C_{1-6} -alk(en/yn)yl, optionally substituted Aryl- C_{3-8} -cycloalk(en)yl, optionally substituted Aryl- C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl, C_{1-6} -alk(en/yn)yl- C_{3-8} -heterocycloalk(en)yl- C_{1-6} -alk(en/yn)yl, C_{3-8} -heterocycloalk(en)yl- C_{1-6} -alk(en/yn)yl, C_{1-6} -alk(en/yn)yl- C_{3-8} -heterocycloalk(en)yl- C_{1-6} -alk(en/yn)yl, Heteroaryl- C_{1-6} -alk(en/yn)yl, Heteroaryl-

- C_{3-8} -cycloalk(en)yl, Heteroaryl- C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl, NR^4R^5 - C_{1-6} -alk(en/yn)yl, NR^4R^5 - C_{3-8} -cycloalk(en)yl, NR^4R^5 - C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl, C_{1-6} -alk(en/yn)yoxy- C_{1-6} -alk(en/yn)yl, C_{3-8} -cycloalk(en)yoxy- C_{1-6} -alk(en/yn)yl, C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yoxy- C_{1-6} -alk(en/yn)yl, halo- C_{1-6} -alk(en/yn)yl, halo- C_{3-8} -cycloalk(en)yl and halo- C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl; wherein each of R^4 and R^5 is independently selected from the group consisting of hydrogen, C_{1-6} -alk(en/yn)yl, C_{3-8} -cycloalk(en)yl and C_{3-8} -cycloalk(en)yl- C_{1-6} -alk(en/yn)yl; as the free base or salts thereof.

10


The compounds of the invention of the general formula I, wherein R^1 , R^2 , R^3 and q are as defined above may be prepared by the methods as represented in the schemes and as described below.

15

In the compounds of the general formulae I - XV, R^1 , R^2 , R^3 and q are as defined under formula I.

Compounds of the general formulae II, VII, VIII, IX, X, XI and XII are either obtained from commercial sources, or prepared by standard methods known to chemists skilled in the art.

20




- Compounds of the general formula III (scheme 1) may be prepared by reacting compounds of the general formula II with bis-(2-haloethyl)ethers, with or without the addition of bases, such as trialkyl amines, potassium carbonate or lithium-, sodium-, or potassium alcoholates, with or without the addition of catalysts such as sodium iodide, in a suitable solvent, such as dimethyl sulfoxide, *N,N*-dimethylformamide or ethanol, at a suitable temperature, such as room temperature or reflux temperature.

30

Compounds of the general formula IV (scheme 1) may be prepared from compounds of the general formula III, by nitration reactions known to chemists skilled in the art, such as reaction with concentrated nitric acid, sodium nitrite or sodium nitrate, in a suitable solvent, such as glacial acetic acid, acetic anhydride, trifluoroacetic acid, concentrated sulfuric acid or mixtures thereof, at appropriate temperatures, for example as described by P.B.D. de la Mare and J.H. Ridd, "Preparative methods of nitration" in *Aromatic substitutions*, pp. 48-56, Butterworths Scientific Publications, London, 1959.


Scheme 2.



Compounds of the general formula V (scheme 2) may be prepared from compounds of the general formula II by nitration reactions known to chemists skilled in the art as described under scheme 1 for the preparation of compounds of the general formula IV.


Compounds of the general formula IV (scheme 2) may be prepared by reacting compounds of the general formula V with suitably substituted bis-(2-haloethyl)ethers as described under scheme 1 for the preparation of compounds of the general formula III.

Scheme 3.




Compounds of the general formula VI (scheme 3) may be prepared from compounds of the general formula IV, by reducing the nitro group to an amino group, with suitable reducing agents such as zinc or iron powder in the presence of acid such as acetic acid or aqueous hydrochloric acid, or by hydrogen gas or ammonium formate in the presence of a suitable hydrogenation catalyst such as palladium on activated carbon in suitable solvents such as methanol, ethanol, ethyl acetate or tetrahydrofuran, at suitable temperatures or under ultrasonic irradiation. Alternatively, tin (II) chloride or sodium dithionite can be used as reducing agents under conditions well known to chemists skilled in the art.

Compounds of the general formula I (scheme 3) may be prepared by reacting compounds of the general formula VI with suitable electrophilic reagents, such as, but not limited to, suitably substituted carboxylic acid fluorides, carboxylic acid chlorides, carboxylic acid bromides, carboxylic acid iodides, carboxylic acid anhydrides, activated esters, chloroformates, and with or without the addition of bases, such as pyridine, trialkyl amines, potassium carbonate, magnesium oxide or lithium-, sodium-, or potassium alcoholates, in a suitable solvent, such as ethyl acetate, dioxane, tetrahydrofuran, acetonitrile or diethyl ether, at suitable temperatures, such as room temperature or reflux temperature. Activated esters and carboxylic acid anhydrides can be prepared from suitably substituted carboxylic acids under conditions known to chemists skilled in the art, for example as described by F. Albericio and L.A. Carpino, "Coupling reagents and activation" in *Methods in enzymology: Solid-phase peptide synthesis*, pp. 104-126, Academic Press, New York, 1997. Carboxylic acid halides can be prepared from suitably substituted carboxylic acids by activation with reagents such as, but not limited to, thionyl chloride, oxalyl chloride, phosphorus tribromide or phosphorus triiodide under conditions well known to chemists skilled in the art.



20 Compounds of the general formula II (scheme 4) may be prepared by reacting compounds of the general formula VII with sodium amide in a suitable solvent, such as xylene at a suitable temperature such as reflux temperature for example as described by J. Lecocq, *Bull. Soc. Chim. Fr.*, 1950, 188.




25 Compounds of the general formula VII, wherein R² is F, Cl, Br or I (scheme 5), may be prepared from compounds of the general formula VIII, by means of

metallation and subsequent reaction with a suitable electrophile known to chemists skilled in the art, using appropriate bases such as butyllithium or lithium di-*t*-butyl(2,2,6,6-tetramethylpiperidino)zincate with subsequent addition of a suitable electrophile such as fluorine, chlorine, bromine, iodine, carbon tetrabromide or hexachloroethane in a suitable solvent such as heptane or tetrahydrofuran, at suitable temperatures, such as -78°C or room temperature for example as described by P. Mongin and G. Quéguiner, *Tetrahedron*, 2001, 57, 4059.

Compounds of the general formula VII, wherein R^1 is F, Cl, Br or I (scheme 5), may be prepared from compounds of the general formula IX, by means of metallation and subsequent electrophilic aromatic substitution as described above.

Scheme 6.



Compounds of the general formula VII, wherein R^2 is $\text{C}_{1-6}\text{-alk(en/yn)yl}$, $\text{C}_{3-8}\text{-cycloalk(en)yl}$, $\text{C}_{3-8}\text{-cycloalk(en)yl-C}_{1-6}\text{-alk(en/yn)yl}$, halo- $\text{C}_{1-6}\text{-alk(en/yn)yl}$, halo- $\text{C}_{3-8}\text{-cycloalk(en)yl}$ or halo- $\text{C}_{3-8}\text{-cycloalk(en)yl-C}_{1-6}\text{-alk(en/yn)yl}$ (scheme 6), may be prepared from compounds of the general formula X, by means of cross-coupling reactions known to chemists skilled in the art, such as *Negishi* coupling (E.-I. Negishi, A.O. King and N. Okukado, *J.Org.Chem.*, 1977, 42, 1821), *Sonogashira* coupling (K. Sonogashira, Y. Tohda and N. Hagihara, *Tet.Lett.*, 1975, 16, 4467), or other transition metal catalyzed cross-coupling reactions such as copper catalyzed reactions (W. Dohle, D.M. Lindsay and P. Knochel, *Org.Lett.*, 2001, 3, 2871).

Compounds of the general formula VII, wherein R^1 is $\text{C}_{1-6}\text{-alk(en/yn)yl}$, $\text{C}_{3-8}\text{-cycloalk(en)yl}$, $\text{C}_{3-8}\text{-cycloalk(en)yl-C}_{1-6}\text{-alk(en/yn)yl}$, halo- $\text{C}_{1-6}\text{-alk(en/yn)yl}$, halo- $\text{C}_{3-8}\text{-cycloalk(en)yl}$ or halo- $\text{C}_{3-8}\text{-cycloalk(en)yl-C}_{1-6}\text{-alk(en/yn)yl}$ (scheme 6), may be prepared from compounds of the general formula XI, by means of cross-coupling reactions as described above.


Additionally, compounds of the general formula VII, wherein R^2 is cyano (scheme 6), may be prepared from compounds of the general formula X, by means of nickel-catalyzed cyanation reactions known to chemists skilled in the art for example as described by L. Cassar, *J. Organomet. Chem.*, 1973, 54, C57-C58.

5

Compounds of the general formula VII, wherein R^1 is cyano (scheme 6), may be prepared from compounds of the general formula XI, by means of nickel-catalyzed cyanation reactions as described above.

- 10 Compounds of the general formula VII, wherein $R^1 = R^2$ (scheme 6), may be prepared from compounds of the general formula X, wherein $R^1 = R^2 = \text{Br}$, by means of cross-coupling reactions or cyanation reactions as described above.

Scheme 7.



- 15 Furthermore, compounds of general formula XIII (scheme 7), wherein R^1 and R^2 are halogen, can be prepared from 2,4,6-trihalopyridines of general formula XII, wherein R^1 and R^2 are halogen, by nitration reactions known to chemists skilled in the art as described under scheme 1 for the preparation of compounds of the general formula IV.
- 20 Compounds of general formula XIV (scheme 7), wherein R^1 and R^2 are halogen, may be prepared by from compounds of general type XII, wherein R^1 and R^2 are halogen, by reaction with morpholine in a suitable solvent such as dimethyl sulfoxide or *N*-methylpyrrolidinone, and with or without the addition of bases, such as pyridine, trialkyl

amines, potassium carbonate, magnesium oxide, at suitable temperatures, such as room temperature or reflux temperature.

Compounds of general type XV may be prepared from compounds of general type XIII by
5 reaction with morpholine in a suitable solvent such as dimethyl sulfoxide or
N-methylpyrrolidinone, and with or without the addition of bases, such as pyridine, trialkyl
amines, potassium carbonate, magnesium oxide, at suitable temperatures, such as room
temperature or reflux temperature. Additionally, compounds of general type XV may be
prepared from compounds of general type XIV by nitration reactions known to chemists
10 skilled in the art as described under scheme 1 for the preparation of compounds of the
general formula IV.

Furthermore, compounds of general formula IV, wherein R¹ or R² or both R¹ and R² is
cyano (scheme 7), may be prepared from compounds of general formula XV, using
15 cyanation reactions as described above. Compounds of general formula III, wherein R¹ or
R² or both R¹ and R² is cyano, may be prepared from compounds of general formula XIV,
using cyanation reactions as described above.

Compounds of the general formula III, wherein R¹ is C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl,
20 C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl or
halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl (scheme 6), may be prepared from compounds of
the general formula XIV, by means of cross-coupling reactions as described above (scheme
6).

25 Compounds of the general formula III, wherein R² is C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl,
C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl or
halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl (scheme 6), may be prepared from compounds of
the general formula XIV, by means of cross-coupling reactions as described above (scheme
6).

30

Compounds of the general formula IV, wherein R¹ is C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl,
C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl or
halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl (scheme 6), may be prepared from compounds of

the general formula XV, by means of cross-coupling reactions as described above (scheme 6).

Compounds of the general formula IV, wherein R² is C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl or halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl (scheme 6), may be prepared from compounds of the general formula XV, by means of cross-coupling reactions as described above (scheme 6).

Compounds of general formula IV, wherein R¹ or R² or both R¹ and R² is C₁₋₆-alk(en/yn)yoxy, C₃₋₈-cycloalk(en)yoxy or C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yoxy, may be prepared from compounds of general formula XV by reaction with the appropriate lithium-, sodium-, or potassium alcoholates or alcohols in the presence of base such as lithium-, sodium-, or potassium hydroxide, lithium-, sodium-, or potassium hydride, and with or without the addition of a catalyst such as copper sulfate, in a suitable solvent such as dioxane, at suitable temperatures, such as room temperature or reflux temperature.

Compounds of general formula III, wherein R¹ or R² or both R¹ and R² is C₁₋₆-alk(en/yn)yoxy, C₃₋₈-cycloalk(en)yoxy or C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yoxy, may be prepared from compounds of general formula XIV by reaction with the appropriate lithium-, sodium-, or potassium alcoholates or alcohols in the presence of base such as lithium-, sodium-, or potassium hydroxide, lithium-, sodium-, or potassium hydride, and with or without the addition of a catalyst such as copper sulfate, in a suitable solvent such as dioxane, at suitable temperatures, such as room temperature or reflux temperature.

Additionally, for further variation of R¹ and R², compounds containing a methoxy-group, can be demethylated by methods known to chemists skilled in the art, such as treatment with boron tribromide in a suitable solvent, such as dichloromethane, at suitable temperatures, such as 0 °C or room temperature. The resulting phenols can then be alkylated by methods known to chemists skilled in the art. Such methods include: (a) the reaction with electrophiles, such as alkyl chlorides, alkyl bromides, alkyl iodides, carbonic acid chlorides, carbonic acid bromides, or carbonic acid anhydrides in the presence of suitable bases, such as potassium carbonate, in a suitable solvent, such as tetrahydrofuran,

N,N-dimethylformamide, or 1,2-dichloroethane, at suitable temperatures, such as room temperature or reflux temperature; (b) the reaction with alkyl alcohols under conditions known as the *Mitsunobu* reaction (O. Mitsunobu, *Synthesis* 1981, 1).

- 5 Compounds containing functional groups, such as hydroxy groups, not compatible with suggested reaction conditions, can be protected and deprotected by methods known to chemists skilled in the art, for example as described by T.W. Greene and P.G.M. Wuts, *Protective groups in organic synthesis*, 2nd edition, Wiley Interscience, 1991. In particular, hydroxy groups can be protected as, but not limited to, methyl-, *tert*-butyl-, trialkylsilyl-,
10 triarylsilyl-, allyl- or trityl ethers.

- Alkynes prepared by Sonogashira reactions may be reduced to alkenes or alkanes by reduction with hydrogen gas or ammonium formate in the presence of a suitable hydrogenation catalyst such as palladium on activated carbon or platinum on activated
15 carbon in suitable solvents such as methanol, ethanol or tetrahydrofuran, at suitable temperatures for example as described by S. Siegel, "Heterogeneous catalytic hydrogenation of C=C and alkynes" in *Comprehensive Organic Synthesis*, v. 8, pp. 417-442, Pergamon Press, 1991.

20

Preparation of the compounds of the invention

Examples

- Analytical LC-MS data were obtained on a PE Sciex API 150EX instrument equipped with
25 atmospheric pressure photo ionisation and a Shimadzu LC-8A/SLC-10A LC system.
Column: 30 X 4.6 mm Waters Symmetry C18 column with 3.5 µm particle size;
Solvent system: A = water/trifluoroacetic acid (100:0.05) and B =
water/acetonitrile/trifluoroacetic acid (5:95:0.03); Method: Linear gradient elution with 90%
A to 100% B in 4 minutes and with a flow rate of 2 mL/minute. Purity was determined by
30 integration of the UV (254 nm) and ELSD trace. The retention times (*t_R*) are expressed in minutes.

Preparative LC-MS-purification was performed on the same instrument with atmospheric pressure chemical ionisation. Column: 50 X 20 mm YMC ODS-A with 5 µm particle size;

Method: Linear gradient elution with 80% A to 100% B in 7 minutes and with a flow rate of 22.7 mL/minute. Fraction collection was performed by split-flow MS detection.

Analytical LC-MS-TOF (TOF = time of flight) data were obtained on a micromass LCT 4-ways MUX equipped with a Waters 2488/Sedex 754 detector system. Column: 30 X 4.6 mm Waters Symmetry C18 column with 3.5 µm particle size; Solvent system: A = water/trifluoroacetic acid (100:0.05) and B = water/acetonitrile/trifluoroacetic acid (5:95:0.03); Method: Linear gradient elution with 90% A to 100% B in 4 minutes and with a flow rate of 2 mL/minute. Purity was determined by integration of the UV (254 nm) and ELSD trace. The retention times (t_R) are expressed in minutes.

GC-MS data were obtained on a Varian CP 3800 gaschromatograph fitted with a Phenomenex column (Zebron ZB-5, length: 15 metres, internal diameter: 0.25 mm) coupled to a Varian Saturn 2000 iontrap mass spectrometer. Method: Duration 15 minutes, column flow 1.4 mL/minute (carrier gas was helium), oven gradient: 0-1 minute, 60 °C; 1-13 minutes, 60-300 °C; 13-15 minutes, 300 °C.

¹H NMR spectra were recorded at 500.13 MHz on a Bruker Avance DRX500 instrument. Deuterated dimethyl sulfoxide (99.8%D) was used as solvent. Tetramethylsilane was used as internal reference standard. Chemical shift values are expressed in ppm-values relative to tetramethylsilane. The following abbreviations are used for multiplicity of NMR signals: s = singlet, d = doublet, t = triplet, q = quartet, qui = quintet, h = heptet, dd = double doublet, ddd = double double doublet, dt = double triplet, dq = double quartet, tt = triplet of triplets, m = multiplet and b = broad singlet.

Preparation of intermediates

4-(4,6-Dimethyl-pyridin-2-yl)-morpholine.

2-Amino-4,6-dimethylpyridine (50 g), bis(2-chloroethyl)ether (57.5 mL), sodium iodide (6.13 g) and triethylamine (137 mL) were mixed in dry *N,N*-dimethylformamide (1 L) under argon and heated to 150 °C for 16 hours. Water/brine/saturated aqueous sodium bicarbonate (2:1:1, 750 mL) were added to the cooled reaction mixture and it was extracted with ethyl acetate (5x 200 mL). The combined organic phases were concentrated *in vacuo* to app. 500

mL. Water (500 mL) and concentrated aqueous hydrochloric acid (35 mL) were added, the phases separated and the aqueous phase washed with ethyl acetate (200 mL). The aqueous phase was made basic with the addition of concentrated aqueous sodium hydroxide (50 mL) and extracted with isopropyl acetate (5x 200 mL). The organic phases were dried over

- 5 magnesium sulfate and concentrated *in vacuo* to furnish 44.0 g (56% yield) of the title compound as a black oil. The crude product was used without further purification. GC-MS (m/z) 192 (M^+); t_R = 5.60. 1H NMR (500 MHz, DMSO- d_6): 2.08 (s, 3H), 2.26 (s, 3H), 3.39 (m, 4H), 3.68 (m, 4H), 6.05 (s, 1H), 6.44 (s, 1H).

- 10 4-(4,6-Dimethyl-5-nitro-pyridin-2-yl)-morpholine.

To 4-(4,6-dimethyl-pyridin-2-yl)-morpholine (9.4 g) dissolved in trifluoroacetic acid (250 mL) cooled to 0 °C was added sodium nitrite (3.54 g) over 15 minutes and the reaction mixture was then stirred 15 minutes at 0 °C. The reaction mixture was concentrated *in vacuo* to app. 100 mL and the pH adjusted to 11 with concentrated aqueous sodium

- 15 hydroxide (150 mL). Brine (200 mL) was added and the mixture was extracted with diethyl ether (4x 150 mL), the organic phase was dried over magnesium sulfate and concentrated *in vacuo*. The crude product was subjected to flash chromatography (SiO₂, heptane/ethylacetate 4:1) to furnish 2.01 g (17% yield) of the title compound as a yellow solid. GC-MS (m/z) 237 (M^+); t_R = 7.69. 1H NMR (500 MHz, DMSO- d_6): 2.28 (s, 3H), 2.39 (s, 3H), 3.60 (m, 4H), 3.67 (m, 4H), 6.72 (s, 1H).
- 20

2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-ylamine.

- Glacial acetic acid (25 mL) was added slowly to a mixture of zinc dust (2.76 g) and 4-(4,6-dimethyl-5-nitro-pyridin-2-yl)-morpholine (2.01 g) in tetrahydrofuran (100 mL) cooled to 0 °C. The reaction mixture was then stirred for 16 hours at 25 °C, filtered through celite, made basic with 25% aqueous ammonia and extracted with tetrahydrofuran (3x 75 mL). The combined organic phases were dried over magnesium sulfate and concentrated *in vacuo* to furnish 1.76 g (100%) of the title compound as a dark red solid. GC-MS (m/z) 207 (M^+); t_R = 7.27. 1H NMR (500 MHz, DMSO- d_6): 2.07 (s, 3H), 2.20 (s, 3H), 3.16 (m, 4H), 3.67 (m, 4H), 4.10 (b, 2H), 6.38 (s, 1H).
- 25
- 30

4-(4,6-Dichloropyridin-2-yl)-morpholine.

Morpholine (5.0 g) was added to a suspension of 2,4,6-trichloropyridine (10.0 g) and sodium carbonate (5.9 g) in acetonitrile (100 mL). The reaction mixture was then stirred at 70 °C for 16 hours, cooled to ambient temperature, filtered through celite and concentrated *in vacuo*. The crude product was subjected to flash chromatography (SiO₂, heptane/ethylacetate 4:1) to furnish 3.90 g (30% yield) of the title compound as an off-white solid. LC-MS (*m/z*) 323.8 (M⁺); *t_R* = 3.10, (UV, ELSD) 98.5%, 98.9%. ¹H NMR (500 MHz, CDCl₃): 3.50 (m, 4H), 3.80 (m, 4H), 6.45 (s, 1H), 6.67 (s, 1H).

4-(4,6-Dichloro-5-nitropyridin-2-yl)-morpholine.

To a solution of 4-(4,6-dichloropyridin-2-yl)-morpholine (3.90 g) in concentrated sulfuric acid (40 mL) was added potassium nitrate (1.80 g) over 10 minutes. The reaction mixture was stirred for 16 hours at ambient temperature and then poured in to crushed ice (500 g). The reaction mixture was made alkaline with concentrated sodium hydroxide and extracted with ethyl acetate (2x100 mL). The combined organic phases were dried over magnesium sulfate and concentrated *in vacuo*. The crude product was subjected to flash chromatography (SiO₂, heptane/ethylacetate 3:1) to furnish 2.26 g (49% yield) of the title compound as a yellow solid. LC-MS (*m/z*) 278.0 (M⁺); *t_R* = 3.10, (UV, ELSD) 96.5%, 98.8%. ¹H NMR (500 MHz, CDCl₃): 3.62 (m, 4H), 3.80 (m, 4H), 6.50 (s, 1H).

4-(4-Chloro-6-methoxy-5-nitropyridin-2-yl)-morpholine and 4-(6-Chloro-4-methoxy-5-nitropyridin-2-yl)-morpholine.

To a solution of 4-(4,6-dichloro-5-nitropyridin-2-yl)-morpholine (2.02 g) in methanol (15 mL) was added sodium methoxide (0.98 g) and the mixture was heated for 16 hours at 65 °C. After cooling to ambient temperature the reaction mixture was concentrated *in vacuo*. The crude product was subjected to flash chromatography (SiO₂, heptane/ethylacetate 3:1) to furnish 0.89 g (45% yield) of 4-(4-chloro-6-methoxy-5-nitropyridin-2-yl)-morpholine (fast eluting band) and 0.38 g (19%) of 4-(6-chloro-4-methoxy-5-nitropyridin-2-yl)-morpholine (late eluting band), both as yellow solids.

4-(4-chloro-6-methoxy-5-nitropyridin-2-yl)-morpholine: LC-MS (*m/z*) 273 (M⁺); *t_R* = 2.77, (UV, ELSD) 95%, 97 %. ¹H NMR (500 MHz, CDCl₃): 3.60 (m, 4H), 3.80 (m, 4H), 3.96 (s, 1H), 6.17 (s, 1H).

4-(6-chloro-4-methoxy-5-nitropyridin-2-yl)-morpholine: LC-MS (m/z) 273 (M^+); t_R = 2.39, (UV, ELSD) 93%, 95%. 1H NMR (500 MHz, $CDCl_3$): 3.57 (m, 4H), 3.80 (m, 4H), 3.95 (s, 3H), 5.95 (s, 1H).

5 4-Chloro-2-methoxy-6-morpholin-4-ylpyridin-3-ylamine

To a solution of 4-(4-chloro-6-methoxy-5-nitropyridin-2-yl)-morpholine (0.82 g) in concentrated hydrochloric acid (50 mL) was added a solution of stannous dichloride (3.38 g) in concentrated hydrochloric acid (80 mL). The reaction mixture was heated to 75 °C for 1 hour and then poured on to crushed ice (400 g) and extracted with ethyl acetate (2x100 mL). The combined organic phases were dried over magnesium sulfate and concentrated *in vacuo*, to furnish 0.45 g (61% yield) of the title compound as an off-white solid. LC-MS (m/z) 244 (M^+); t_R = 1.48, (UV, ELSD) 89%, 94%. 1H NMR (500 MHz, $CDCl_3$): 3.30 (m, 4H), 3.65 (br s, 2H), 3.85 (m, 4H), 3.97 (s, 3H), 6.20 (s, 1H).

15 2-Chloro-4-methoxy-6-morpholin-4-ylpyridin-3-ylamine

To a solution of 4-(6-chloro-4-methoxy-5-nitropyridin-2-yl)-morpholine (0.38 g) in concentrated hydrochloric acid (20 mL) was added a solution of stannous dichloride (1.57 g) in concentrated hydrochloric acid (60 mL). The reaction mixture was heated to 75 °C for 5 minutes and then poured on to crushed ice (100 g) and extracted with ethyl acetate (2x20 mL). The combined organic phases were dried over magnesium sulfate and concentrated *in vacuo*, to furnish 0.28 g (83% yield) of the title compound as an off-white solid. 1H NMR (500 MHz, $CDCl_3$): 3.35 (m, 4H), 3.65 (br s, 2H), 3.80 (m, 4H), 3.90 (s, 3H), 6.10 (s, 1H).

Compounds of the invention

25 Acid addition salts of the compounds of the invention may easily be formed by methods known to the person skilled in the art.

Example 1

30 *1aa* (2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-carbamic acid benzyl ester.

Benzyl chloroformate (18 mg) was added to a solution of 0.085 M 2,4-dimethyl-6-morpholin-4-yl-pyridin-3-ylamine and 0.17 M *N,N*-diisopropyl-ethylamine in 1,2-dichloroethane (1 mL). The vial was shaken for 16 hours under argon and concentrated *in*

vacuo. Aqueous sodium hydroxide (1 M, 1 mL) was added and the crude mixture was extracted with isopropyl acetate/tetrahydrofuran (4:1, 2x 1 mL). The organic phase was washed with brine (1 mL), concentrated *in vacuo* and redissolved in 1-propanol/dimethyl sulfoxide (1:1, 0.4 mL) of which 0.2 mL was subjected to preparative LC-MS purification to furnish 4.5 mg (31% yield) of the title compound as an oil. LC-MS (m/z) 342 (MH^+); t_R = 1.58, (UV, ELSD) 99%, 99%.

The following compounds were prepared analogously:

Iab (2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-carbamic acid 2-chloro-benzyl ester.

10 Yield: 18%. LC-MS (m/z) 376 (MH^+); t_R = 1.78, (UV, ELSD) 99%, 100%.

Iac 2-(4-Chloro-phenyl)-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide.

Yield: 4%. LC-MS (m/z) 360 (MH^+); t_R = 1.59, (UV, ELSD) 96%, 100%.

15 *Iad* 2-Phenyl-cyclopropanecarboxylic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide.

Yield: 24%. LC-MS (m/z) 352 (MH^+); t_R = 1.64, (UV, ELSD) 96%, 100%.

Iae N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-thiophen-2-yl-acetamide.

20 Yield: 16%. LC-MS (m/z) 332 (MH^+); t_R = 1.20, (UV, ELSD) 93%, 99%.

Iaf 3-Cyclohexyl-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-propionamide.

Yield: 15%. LC-MS (m/z) 346 (MH^+); t_R = 1.81, (UV, ELSD) 91%, 100%.

25 *Iag* (2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-carbamic acid isobutyl ester.

Yield: 29%. LC-MS (m/z) 308 (MH^+); t_R = 1.44, (UV, ELSD) 97%, 99%.

Iah 3-(3-Chloro-phenyl)-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-propionamide.

30 3-(3-Chlorophenyl)propionic acid (20 mg) was stirred at 25°C for 2 hours under argon in oxalyl chloride (2 M in dichloromethane, 1 mL). The solvent was removed *in vacuo* and a solution of 0.085 M 2,4-dimethyl-6-morpholin-4-yl-pyridin-3-ylamine and 0.17 M *N,N*-diisopropyl-ethylamine in 1,2-dichloroethane (1 mL) was added to the reaction mixture. The vial was shaken for 16 hours under argon and concentrated *in vacuo*. Aqueous sodium

hydroxide (1 M, 1 mL) was added and the crude mixture was extracted with isopropyl acetate/tetrahydrofuran (4:1, 2x 1 mL). The organic phase was washed with brine (1 mL), concentrated *in vacuo* and redissolved in 1-propanol/dimethyl sulfoxide (1:1, 0.4 mL) of which 0.2 mL was subjected to preparative LC-MS purification to furnish 2.3 mg (14%
5 yield) of the title compound as an oil. LC-MS (m/z) 374 (MH^+); t_R = 1.71, (UV, ELSD) 99%, 99%.

The following compounds were prepared analogously:

Iai N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(3,5-dimethyl-phenyl)-acetamide.

10 Yield: 19%. LC-MS (m/z) 354 (MH^+); t_R = 1.69, (UV, ELSD) 99%, 99%.

Iaj N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-3-p-tolyl-propionamide.

Yield: 20%. LC-MS (m/z) 354 (MH^+); t_R = 1.64, (UV, ELSD) 99%, 100%.

15 *Iak 2-(3-Chloro-phenyl)-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide.*

Yield: 14%. LC-MS (m/z) 360 (MH^+); t_R = 1.58, (UV, ELSD) 97%, 99%.

Ial 2-(3,4-Dichloro-phenyl)-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide.

Yield: 9%. LC-MS (m/z) 395 (MH^+); t_R = 1.84, (UV, ELSD) 97%, 99%.

20

Iam N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-thiophen-3-yl-acetamide.

Yield: 18%. LC-MS (m/z) 332 (MH^+); t_R = 1.18, (UV, ELSD) 97%, 99%.

Ian N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-p-tolyl-acetamide.

25 Yield: 16%. LC-MS (m/z) 340 (MH^+); t_R = 1.50, (UV, ELSD) 96%, 99%.

Iao 2-(3-Bromo-phenyl)-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide.

Yield: 12%. LC-MS (m/z) 405 (MH^+); t_R = 1.63, (UV, ELSD) 96%, 99%.

30 *Iap N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(3-trifluoromethyl-phenyl)-acetamide.*

Yield: 20%. LC-MS (m/z) 394 (MH^+); t_R = 1.77, (UV, ELSD) 94%, 99%.

Iaq N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-phenyl-acetamide.

Yield: 11%. LC-MS (m/z) 326 (MH^+); t_R = 1.29, (UV, ELSD) 93%, 99%.

Iar 3,5,5-Trimethyl-hexanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide.

5 Yield: 20%. LC-MS (m/z) 348 (MH^+); t_R = 1.97, (UV, ELSD) 93%, 99%.

Ias Octanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide.

Yield: 44%. LC-MS (m/z) 334 (MH^+); t_R = 1.92, (UV, ELSD) 92%, 99%.

10 *Iat N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-naphthalen-2-yl-acetamide.*

Yield: 4%. LC-MS (m/z) 376 (MH^+); t_R = 1.73, (UV, ELSD) 92%, 99%.

Iau Heptanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide.

Yield: 24%. LC-MS (m/z) 320 (MH^+); t_R = 1.56, (UV, ELSD) 90%, 99%.

15

Iav N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(3,4-dimethyl-phenyl)-acetamide.

Yield: 26%. LC-MS (m/z) 354 (MH^+); t_R = 1.65, (UV, ELSD) 77%, 99%.

Iaw 2-Cyclohex-1-enyl-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide.

20 Yield: 13%. LC-MS (m/z) 330 (MH^+); t_R = 1.50, (UV, ELSD) 72%, 99%.

Iax N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(4-methoxy-3-methyl-phenyl)-acetamide.

Yield: 16%. LC-MS (m/z) 370 (MH^+); t_R = 1.56, (UV, ELSD) 94%, 99%.

25

Iay N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(4-methoxy-phenyl)-acetamide.

Yield: 19%. LC-MS (m/z) 356 (MH^+); t_R = 1.35, (UV, ELSD) 96%, 99%.

Iaz N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-3-(4-methoxy-phenyl)-propionamide.

30 Yield: 15%. LC-MS (m/z) 370 (MH^+); t_R = 1.48, (UV, ELSD) 76%, 99%.

1ba N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-m-tolyl-acetamide.

m-Tolylacetic acid (0.33 g), *N,N*-diisopropyl-ethylamine (0.90 mL) and *N*-[(dimethylamino)-1*H*-1,2,3-triazolo-[4,5-*b*]pyridin-1-yl-methylene]-*N*-methyl-methanaminium hexafluoro-phosphate *N*-oxide (1.00 g) were mixed in dry

- 5 *N,N*-dimethylformamide (3 mL) and stirred under argon for 2 minutes. 2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-ylamine (0.30 g) dissolved in dry *N,N*-dimethylformamide (2 mL) was added to the reaction mixture, which was stirred at 25 °C under argon for 16 hours. Ethyl acetate (20 mL) was added and the organic phase was washed with saturated aqueous ammonium chloride/water (1:1, 20 mL), water (20 mL), brine/water (1:1, 20 mL), dried
10 over sodium sulfate, concentrated *in vacuo* and purified by flash chromatography (SiO₂, heptane/ethylacetate 3:1) to furnish 0.069 g (14% yield) of the title compound as a white solid. LC-MS (*m/z*) 340 (MH⁺); *t_R* = 1.42, (UV, ELSD) 96%, 100%. ¹H NMR (500 MHz, DMSO-*d*₆): 2.00 (s, 3H), 2.11 (s, 3H), 2.29 (s, 3H), 3.37 (m, 4H), 3.56 (s, 2H), 3.67 (m, 4H), 6.52 (s, 1H), 7.06 (d, 1H), 7.15 (m, 2H), 7.21 (t, 1H), 9.30 (s, 1H).

15

The following compounds were prepared analogously:

1bb N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(4-fluoro-phenyl)-acetamide.

- Yield: 14%. LC-MS (*m/z*) 344 (MH⁺); *t_R* = 1.34, (UV, ELSD) 99%, 99%. ¹H NMR (500 MHz, DMSO-*d*₆): 1.99 (s, 3H), 2.10 (s, 3H), 3.37 (m, 4H), 3.60 (s, 2H), 3.66 (m, 4H), 6.52
20 (s, 1H), 7.16 (dd, 2H), 7.38 (dd, 2H), 9.33 (s, 1H).

1bc N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-3,3-dimethyl-butyramide.

- Yield: 53%. LC-MS (*m/z*) 306 (MH⁺); *t_R* = 1.26, (UV, ELSD) 99%, 98%. ¹H NMR (500 MHz, DMSO-*d*₆): 1.05 (s, 9H), 2.07 (s, 3H), 2.18 (s, 2H), 2.19 (s, 3H), 3.37 (m, 4H), 3.67
25 (m, 4H), 6.54 (s, 1H), 9.01 (s, 1H).

1bd N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(3-fluoro-phenyl)-acetamide.

- Yield: 15%. LC-MS (*m/z*) 344 (MH⁺); *t_R* = 1.54, (UV, ELSD) 100%, 100%. ¹H NMR (500 MHz, DMSO-*d*₆): 2.00 (s, 3H), 2.11 (s, 3H), 3.37 (m, 4H), 3.64 (s, 2H), 3.66 (m, 4H), 6.52
30 (s, 1H), 7.08 (dt, 1H), 7.18 (m, 2H), 7.38 (m, 1H), 9.34 (s, 1H).

1be 2-Bicyclo[2.2.1]hept-2-yl-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide.

Yield: 62%. LC-MS (m/z) 344 (MH^+); t_R = 1.58, (UV, ELSD) 99%, 99%. 1H NMR (500 MHz, DMSO- d_6): 1.14 (m, 4H), 1.42 (m, 4H), 1.90 (m, 1H), 2.01 (m, 1H), 2.04 (s, 3H),
5 2.10 (m, 1H), 2.16 (s, 3H), 2.21 (m, 2H), 3.37 (m, 4H), 3.67 (m, 4H), 6.53 (s, 1H), 9.04 (s, 1H).

1bf 2-(3,4-Difluoro-phenyl)-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide.

Yield: 9%. LC-MS (m/z) 362 (MH^+); t_R = 1.52, (UV, ELSD) 95%, 99%. 1H NMR (500
10 MHz, DMSO- d_6): 2.00 (s, 3H), 2.11 (s, 3H), 3.37 (m, 4H), 3.63 (s, 2H), 3.66 (m, 4H), 6.52 (s, 1H), 7.19 (m, 1H), 7.39 (m, 2H), 9.32 (s, 1H).

1bg 4-Methyl-pentanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide.

Yield: 34%. LC-MS (m/z) 306 (MH^+); t_R = 1.33, (UV, ELSD) 100%, 99%. 1H NMR (500
15 MHz, DMSO- d_6): 0.91 (d, 6H), 1.49 (dt, 2H), 1.58 (m, 1H), 2.04 (s, 3H), 2.16 (s, 3H), 2.28 (t, 2H), 3.37 (m, 4H), 3.67 (m, 4H), 6.53 (s, 1H), 9.07 (s, 1H).

1bh 2-Cyclopent-2-enyl-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide.

Yield: 13%. LC-MS (m/z) 316 (MH^+); t_R = 1.25, (UV, ELSD) 97%, 94%. 1H NMR (500
20 MHz, DMSO- d_6): 1.51 (m, 1H), 2.05 (m, 1H), 2.06 (s, 3H), 2.17 (s, 3H), 2.26 (m, 2H), 2.35 (m, 2H), 3.07 (m, 1H), 3.38 (m, 4H), 3.68 (m, 4H), 5.73 (m, 1H), 5.77 (m, 1H), 6.54 (s, 1H), 9.09 (s, 1H).

1bi 2-Cyclohexyl-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide.

25 Yield: 12%. LC-MS (m/z) 332 (MH^+); t_R = 1.50, (UV, ELSD) 99%, 95%. 1H NMR (500 MHz, DMSO- d_6): 0.98 (m, 2H), 1.20 (m, 3H), 1.71 (m, 6H), 2.05 (s, 3H), 2.15 (d, 2H), 2.16 (s, 3H), 3.37 (m, 4H), 3.67 (m, 4H), 6.53 (s, 1H), 9.05 (s, 1H).

1bj 5-Methyl-hexanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide.

30 Yield: 40%. LC-MS-TOF (m/z) 320 (MH^+); t_R = 1.51, (UV, ELSD) 97%, 100%. 1H NMR (500 MHz, DMSO- d_6): 0.87 (d, 6H), 1.21 (m, 2H), 1.60 (m, 3H), 2.05 (s, 3H), 2.16 (s, 3H), 2.25 (t, 2H), 3.37 (m, 4H), 3.67 (m, 4H), 6.53 (s, 1H), 9.05 (s, 1H).

1bk 2-Cyclopentyl-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide.

2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-ylamine (0.22 g) and cyclopentylacetyl chloride (0.19 mL) were dissolved in acetonitrile (5 mL) and heated to 150 °C for 10 minutes in a sealed microwave process vial. The reaction mixture was concentrated *in vacuo* and purified
5 by flash chromatography (SiO₂, heptane/ethylacetate 3:1) to furnish 0.17 g (49% yield) of the title compound as a white solid. LC-MS (*m/z*) 318 (MH⁺); *t_R* = 1.40, (UV, ELSD) 97%, 99%. ¹H NMR (500 MHz, DMSO-*d*₆): 1.21 (m, 2H), 1.52 (m, 2H), 1.61 (m, 2H), 1.77 (m, 2H), 2.05 (s, 3H), 2.17 (s, 3H), 2.24 (m, 1H), 2.26 (m, 2H), 3.37 (m, 4H), 3.67 (m, 4H), 6.53 (s, 1H), 9.05 (s, 1H).

10

The following compounds were prepared analogously except *1bl* and *1bm* which were recrystallized from ethyl acetate after flash chromatography:

1bl 3-Cyclopentyl-N-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-propionamide.

Yield: 34%. LC-MS (*m/z*) 332 (MH⁺); *t_R* = 1.57, (UV, ELSD) 99%, 99%. ¹H NMR (500
15 MHz, DMSO-*d*₆): 1.11 (m, 2H), 1.49 (m, 2H), 1.60 (m, 4H), 1.77 (m, 3H), 2.04 (s, 3H), 2.16 (s, 3H), 2.28 (t, 2H), 3.37 (m, 4H), 3.67 (m, 4H), 6.53 (s, 1H), 9.06 (s, 1H).

1bm Hexanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide.

Yield: 51%. LC-MS (*m/z*) 306 (MH⁺); *t_R* = 1.39, (UV, ELSD) 99%, 99%. ¹H NMR (500
20 MHz, DMSO-*d*₆): 0.88 (t, 3H), 1.31 (m, 4H), 1.60 (m, 2H), 2.05 (s, 3H), 2.16 (s, 3H), 2.27 (t, 2H), 3.37 (m, 4H), 3.67 (m, 4H), 6.53 (s, 1H), 9.03 (s, 1H).

1bn N-(4-Chloro-2-methoxy-6-morpholin-4-yl-pyridin-3-yl)-2-cyclopentylacetamide.

Yield: 53%. LC-MS (*m/z*) 354 (MH⁺); *t_R* = 2.68, (UV, ELSD) 98%, 99%. ¹H NMR (500
25 MHz, CDCl₃): 1.25 (m, 2H), 1.50-1.65 (m, 4H), 1.90 (m, 2H), 2.45 (m, 3H), 3.45 (m, 4H), 3.77 (m, 4H), 3.90 (s, 3H), 6.20 (s, 1H), 6.50 (s, 1H).

1bo N-(2-Chloro-4-methoxy-6-morpholin-4-yl-pyridin-3-yl)-2-cyclopentylacetamide.

Yield: 69%. LC-MS (*m/z*) 354 (MH⁺); *t_R* = 2.39, (UV, ELSD) 99%, 99%. ¹H NMR (500
30 MHz, CDCl₃): 1.25 (m, 2H), 1.50-1.70 (m, 4H), 1.90 (m, 2H), 2.35 (m, 3H), 3.50 (m, 4H), 3.80 (m, 4H), 3.85 (s, 3H), 6.00 (s, 1H), 6.45 (s, 1H).

1bp N-(2-Chloro-4-methoxy-6-morpholin-4-yl-pyridin-3-yl)-3,3-dimethylbutyramide.

Yield: 56%. LC-MS (m/z) 342 (MH^+); t_R = 2.31, (UV, ELSD) 99%, 99%. 1H NMR (500 MHz, $CDCl_3$): 1.10 (s, 9H), 2.25 (s, 2H), 3.50 (m, 4H), 3.77 (m, 4H), 3.85 (s, 3H), 6.00 (s, 1H), 6.45 (s, 1H).

5

1bq N-(4-Chloro-2-methoxy-6-morpholin-4-yl-pyridin-3-yl)-3,3-dimethylbutyramide.

Yield: 68%. LC-MS (m/z) 342 (MH^+); t_R = 1.39, (UV, ELSD) 99%, 99%. 1H NMR (500 MHz, $DMSO-d_6$): 1.10 (s, 9H), 2.15 (s, 2H), 3.45 (m, 4H), 3.70 (m, 4H), 3.80 (s, 3H), 6.45 (s, 1H), 8.95 (s, 1H).

10

1br N-(4-Chloro-2-methoxy-6-morpholin-4-yl-pyridin-3-yl)-propionamide.

Yield: 71%. LC-MS (m/z) 300 (MH^+); t_R = 0.97, (UV, ELSD) 98%, 98%. 1H NMR (500 MHz, $DMSO-d_6$): 1.05 (t, 3H), 2.25 (q, 2H), 3.45 (m, 4H), 3.70 (m, 4H), 3.80 (s, 3H), 6.45 (s, 1H), 9.00 (s, 1H).

15

Table 1. Reagents used for the preparation of compounds in Example 1.

Name	Supplier	CAS no.	Cat.no.
1-Cyclohexenylacetic acid	Alfa	18294-87-6	19462
3,4-Difluorophenylacetic acid	ABCR	658-93-5	F02874E
3-Bromophenylacetic acid	Aldrich	1878-67-7	28,886-1
3-Chlorophenylacetic acid	Aldrich	1878-65-5	C6,335-9
3-(Trifluoromethyl)phenylacetic acid	Aldrich	351-35-9	19,335-6
2-Amino-4,6-dimethylpyridine	Aldrich	5407-87-4	A5,180-7
2-Chlorobenzyl chloroformate	Aldrich	39545-31-8	49,379-1
2-Cyclopentene-1-acetic acid	Aldrich	13668-61-6	C11,285-2
2-Naphthylacetic acid	Aldrich	581-96-4	31,791-8
2-Phenylacetic acid	Aldrich	103-82-2	P1,662-1
2,4,6-trichloropyridine	Aldrich	18083-89-7	63,353-4
3-(3-Chlorophenyl)propionic acid	ABCR	21640-48-2	TWC2925
3-(4-Methoxyphenyl)propionic acid	Aldrich	1929-29-9	M2,352-7
3-(4-Methylphenyl)propionic acid	Aldrich	1505-50-6	11,826-5
3,4-Dichlorophenylacetic acid	Aldrich	5807-30-7	28,000-3

3,4-Dimethylphenylacetic acid	Vitas-M	17283-16-8	TBB000367
3,5,5-Trimethylhexanoic acid	Acros	3302-10-1	26944-0250
3,5-Dimethylphenylacetic acid	ABCR	42288-46-0	C-42288-46
3-Cyclohexylpropionyl chloride	Acros	39098-75-4	35071-0250
3-Cyclopentylpropionyl chloride	Aldrich	104-97-2	26,859-3
3-Fluorophenylacetic acid	Aldrich	331-25-9	24,804-5
4-Chlorophenylacetyl chloride	Lancaster	25026-34-0	6317
4-Fluorophenylacetic acid	Aldrich	405-50-5	F1,330-4
4-Methoxy-3-methylphenylacetic acid	Vitas-M	4513-73-9	TBB000371
4-Methoxyphenylacetic acid	Aldrich	104-01-8	M1,920-1
4-Methylpentanoic acid	Aldrich	646-07-1	27,782-7
5-Methylhexanoic acid	Matrix	628-46-6	3527
Benzyl chloroformate	Aldrich	501-53-1	11,993-8
Bicyclo[2.2.1]hept-2-yl-acetic acid	Aldrich	1007-01-8	12,726-4
Bis-(2-chloroethyl)ether	Aldrich	111-44-4	C4,113-4
Cyclohexyl-acetic acid	Aldrich	5292-21-7	C10,450-7
Cyclopentylacetyl chloride	Lancaster	1122-99-2	14562
Heptanoic acid	Aldrich	111-14-8	14,687-0
Hexanoyl chloride	Aldrich	142-61-0	29,465-9
Isobutyl chloroformate	Aldrich	543-27-1	17,798-9
<i>m</i> -Tolylacetic acid	Aldrich	621-36-3	T3,809-1
<i>N</i> -[(Dimethylamino)- <i>1H</i> -1,2,3-triazolo[4,5- <i>b</i>] pyridin-1-ylmethylene]- <i>N</i> -methylmethanaminium hexafluorophosphate <i>N</i> -oxide	Fluka	148893-10-1	11373
Octanoic acid	Aldrich	124-07-2	15,375-3
Oxalyl chloride	Aldrich	79-37-8	32,042-0
<i>p</i> -Tolylacetic acid	Aldrich	622-47-9	T3,810-5
sodium iodide	Aldrich	7681-82-5	32,245-8
sodium nitrite	Aldrich	7632-00-0	51,091-2
<i>Tert</i> -butylacetic acid	Aldrich	1070-83-3	B8,840-3

Thiophen-2-acetyl chloride	Aldrich	39098-97-0	19,599-5
Thiophene-3-acetic acid	Aldrich	6964-21-2	22,063-9
Trans-2-phenyl-1-cyclopropanecarbonyl chloride	Aldrich	939-87-7	13,430-9
Zinc	Aldrich	52374-36-4	20,998-8

In vitro and in vivo testing

The compounds of the invention have been tested and shown effect in at least one of the
5 below models:

Relative efflux through the KCNQ2 channel.

This exemplifies a KCNQ2 screening protocol for evaluating compounds of the present invention. The assay measures the relative efflux through the KCNQ2 channel, and was
10 carried out according to a method described by Tang et al. (Tang, W. et. al., J. Biomol. Screen. 2001, 6, 325-331) for hERG potassium channels with the modifications described below.

An adequate number of CHO cells stably expressing voltage-gated KCNQ2 channels were
15 plated at a density sufficient to yield a confluent mono-layer on the day before the experiment. The cells were loaded with 1 $\mu\text{Ci/ml}$ [^{86}Rb] over night. On the day of the experiment cells were washed with a HBSS-containing buffer (Hanks balanced salt solution provided from Invitrogen, cat# 14025-050). Cells were pre-incubated with drug for 30 minutes and the $^{86}\text{Rb}^+$ efflux was stimulated by a submaximal concentration of 15 mM
20 potassium chloride in the continued presence of drug for additional 30 minutes. After a suitable incubation period, the supernatant was removed and counted in a liquid scintillation counter (Tricarb). Cells were lysed with 2 mM sodium hydroxide and the amount of $^{86}\text{Rb}^+$ was counted. The relative efflux was calculated $((\text{CPM}_{\text{super}}/(\text{CPM}_{\text{super}} + \text{CPM}_{\text{cell}}))_{\text{Cmpd}}/(\text{CPM}_{\text{super}}/(\text{CPM}_{\text{super}} + \text{CPM}_{\text{cell}}))_{15\text{mM KCl}}) * 100 - 100$.

25

The compounds of the invention have an EC_{50} of less than 20000nM, in most cases less than 2000 nM and in many cases less than 200 nM. Accordingly, the compounds of the invention

are considered to be useful in the treatment of diseases associated with the KCNQ family potassium channels.

Electrophysiological patch-clamp recordings in CHO cells

- 5 Voltage-activated KCNQ2 currents were recorded from mammalian CHO cells by use of conventional patch-clamp recordings techniques in the whole-cell patch-clamp configuration (Hamill OP et.al. *Pflügers Arch* 1981; 391: 85-100). CHO cells with stable expression of voltage-activated KCNQ2 channels were grown under normal cell culture conditions in CO₂ incubators and used for electrophysiological recordings 1-7 days after
- 10 plating. KCNQ2 potassium channels were activated by voltage steps up to + 80 mV in increments of 5-20 mV (or with a ramp protocol) from a membrane holding potential between - 100 mV and - 40 mV (Tatulian L et al. *J Neuroscience* 2001; 21 (15): 5535-5545). The electrophysiological effects induced by the compounds were evaluated on various parameters of the voltage-activated KCNQ2 current. Especially effects on the
- 15 activation threshold for the current and on the maximum induced current were studied.

- Some of the compounds of the invention have been tested in this test. A left-ward shift of the activation threshold or an increase in the maximum induced potassium current is expected to decrease the activity in neuronal networks and thus make the compounds useful
- 20 in diseases with increased neuronal activity - like epilepsy.

Electrophysiological recordings of KCNQ2, KCNQ2/KCNQ3 or KCNQ5 channels in oocytes

- Voltage-activated KCNQ2, KCNQ2/KCNQ3 or KCNQ5 currents were recorded from
- 25 *Xenopus* oocytes injected with mRNA coding for KCNQ2, KCNQ2+KCNQ3 or KCNQ5 ion channels (Wang et al., *Science* 1998, 282, 1890-1893; Lerche et al., *J Biol Chem* 2000, 275, 22395-400). KCNQ2, KCNQ2/KCNQ3 or KCNQ5 potassium channels were activated by voltage steps from the membrane holding potential (between - 100 mV and - 40 mV) up to + 40 mV in increments of 5-20 mV (or by a ramp protocol). The electrophysiological
- 30 effects induced by the compounds were evaluated on various parameters of the voltage-activated KCNQ2, KCNQ2/KCNQ3 or KCNQ5 currents. Especially effects on the activation threshold for the current and on the maximum induced current were studied.

The hyperpolarizing effects of some of the compounds were also tested directly on the membrane potential during current clamp.

Maximum electroshock

- 5 The test was conducted in groups of male mice using corneal electrodes and administering a square wave current of 26mA for 0.4 seconds in order to induce a convulsion characterised by a tonic hind limb extension (Wlaz et al. *Epilepsy Research* 1998, 30, 219-229).

Pilocarpine induced seizures

- 10 Pilocarpine induced seizures are induced by intraperitoneal injection of pilocarpine 250mg/kg to groups of male mice and observing for seizure activity resulting in loss of posture within a period of 30 minutes (Starr et al. *Pharmacology Biochemistry and Behavior* 1993, 45, 321-325).

Electrical seizure -threshold test

- A modification of the up-and-down method (Kimball et al. *Radiation Research* 1957, 1-12) was used to determine the median threshold to induce tonic hind-limb extension in response to corneal electroshock in groups of male mice. The first mouse of each group received an electroshock at 14 mA, (0.4 s, 50 Hz) and was observed for seizure activity. If a seizure was
20 observed the current was reduced by 1 mA for the next mouse, however, if no seizure was observed then the current was increased by 1 mA. This procedure was repeated for all 15 mice in the treatment group.

Chemical seizure -threshold test

- 25 The threshold dose of pentylenetetrazole required to induce a clonic convulsion was measured by timed infusion of pentylenetetrazole (5mg / mL at 0.5 mL/minute) into a lateral tail vein of groups of male mice (Nutt et al. *J Pharmacy and Pharmacology* 1986, 38, 697-698).

30 Amygdala kindling

Rats underwent surgery to implantation of tri-polar electrodes into the dorsolateral amygdala. After surgery the animals were allowed to recover before the groups of rats received either varying doses of test compound or the drug's vehicle. The animals were

stimulated with their initial after discharge threshold + 25 μ A daily for 3-5 weeks and on each occasion seizure severity, seizure duration, and duration of electrical after discharge were noted. (Racine. *Electroencephalography and Clinical Neurophysiology* 1972, 32, 281-294).

5

Side effects

Central nervous system side-effects were measured by measuring the time mice would remain on rotarod apparatus (Capacio et al. *Drug and Chemical Toxicology* 1992, 15, 177-201); or by measuring their locomotor activity by counting the number of infra-red beams crossed in a test cage (Watson et al. *Neuropharmacology* 1997, 36, 1369-1375).

10


Hypothermic actions on the animals core body temperature of the compound were measured by either rectal probe or implanted radiotelemetry transmitters capable of measuring temperature (Keeney et al. *Physiology and Behaviour* 2001, 74, 177-184).

15 Pharmacokinetics

The pharmacokinetic properties of the compounds were determined via. i.v. and p.o. dosing to Sprague Dawley rats, and, thereafter, drawing blood samples over 20 hours. Plasma concentrations were determined with LC/MS/MS.

CLAIMS

1. A compound having the general formula I:



wherein:

5 q is 0 or 1;

each of R¹ and R² is independently selected from the group consisting of halogen, cyano, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl, halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yloxy, C₃₋₈-cycloalk(en)yloxy and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yloxy; and

10 R³ is selected from the group consisting of C₁₋₈-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, optionally substituted Aryl-C₁₋₆-alk(en/yn)yl, optionally substituted Aryl-C₃₋₈-cycloalk(en)yl, optionally substituted Aryl-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yl-C₃₋₈-heterocycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₃₋₈-heterocycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yl-C₃₋₈-heterocycloalk(en)yl-C₁₋₆-alk(en/yn)yl, Heteroaryl-C₁₋₆-alk(en/yn)yl, Heteroaryl-C₃₋₈-cycloalk(en)yl, Heteroaryl-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, NR⁴R⁵-C₁₋₆-alk(en/yn)yl, NR⁴R⁵-C₃₋₈-cycloalk(en)yl, NR⁴R⁵-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yloxy-C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yloxy-C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yloxy-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl and halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl; wherein

20 each of R⁴ and R⁵ is independently selected from the group consisting of hydrogen, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl; as the free base or salts thereof.


25

2. A compound according to claim 1 wherein q is 0.
3. A compound according to claim 1 wherein q is 1.
- 5 4. A compound according to any one of claims 1-3 wherein each of R¹ and R² is independently selected from the group consisting of C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yoxy and halogen.
- 10 5. A compound according to claim 4 wherein both R¹ and R² are C₁₋₆-alk(en/yn)yl.
6. A compound according to claim 4 wherein R¹ is C₁₋₆-alk(en/yn)yoxy and R² is halogen, or wherein R¹ is halogen and R² is C₁₋₆-alk(en/yn)yoxy.
- 15 7. A compound according to any one of claims 1-6 wherein R³ is selected from the group consisting of C₁₋₈-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, optionally substituted Aryl-C₁₋₆-alk(en/yn)yl, optionally substituted Aryl-C₃₋₈-cycloalk(en)yl, optionally substituted Aryl-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, Heteroaryl-C₁₋₆-alk(en/yn)yl, Heteroaryl-C₃₋₈-cycloalk(en)yl, Heteroaryl-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl,
- 20 8. A compound according to claim 7 wherein R³ is selected from the group consisting of C₁₋₈-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, optionally substituted Aryl-C₁₋₆-alk(en/yn)yl, optionally substituted Aryl-C₃₋₈-cycloalk(en)yl and Heteroaryl-C₁₋₆-alk(en/yn)yl.
- 25 9. A compound according to any one of claims 1, 7 and 8 wherein optionally substituted Aryl may be substituted with one or more substituent indepently selected from the group consisting of halogen, cyano, C₁₋₆-alk(en/yn)yl, C₃₋₈-cycloalk(en)yl, C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl, halo-C₃₋₈-cycloalk(en)yl, halo-C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yl, C₁₋₆-alk(en/yn)yoxy, C₃₋₈-cycloalk(en)yoxy and C₃₋₈-cycloalk(en)yl-C₁₋₆-alk(en/yn)yoxy.
- 30

10. A compound according to claim 9 wherein optionally substituted Aryl may be substituted with one or more substituent indepently selected from the group consisting of halogen, C₁₋₆-alk(en/yn)yl, halo-C₁₋₆-alk(en/yn)yl and C₁₋₆-alk(en/yn)yoxy.
- 5 11. A compound according to any one of Claims 1-10, said compounds being selected from the group consisting of:
- (2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-carbamic acid benzyl ester;
 - (2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-carbamic acid 2-chloro-benzyl ester;
 - 2-(4-Chloro-phenyl)-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide;
 - 2-Phenyl-cyclopropanecarboxylic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide;
 - N*-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-thiophen-2-yl-acetamide;
 - 3-Cyclohexyl-*N*-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-propionamide;
 - (2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-carbamic acid isobutyl ester;
 - 3-(3-Chloro-phenyl)-*N*-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-propionamide;
 - N*-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(3,5-dimethyl-phenyl)-acetamide;
 - N*-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-3-*p*-tolyl-propionamide;
 - 2-(3-Chloro-phenyl)-*N*-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide;
 - 2-(3,4-Dichloro-phenyl)-*N*-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide;
 - N*-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-thiophen-3-yl-acetamide;
 - N*-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-*p*-tolyl-acetamide;
 - 2-(3-Bromo-phenyl)-*N*-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide;
 - N*-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(3-trifluoromethyl-phenyl)-acetamide;
 - N*-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-phenyl-acetamide;
 - 3,5,5-Trimethyl-hexanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide;
 - Octanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide;
 - N*-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-naphthalen-2-yl-acetamide;
 - Heptanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide;
 - N*-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(3,4-dimethyl-phenyl)-acetamide;
 - 2-Cyclohex-1-enyl-*N*-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide;
 - N*-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(4-methoxy-3-methyl-phenyl)-

acetamide;
N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(4-methoxy-phenyl)-acetamide;
N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-3-(4-methoxy-phenyl)-propionamide;
N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-*m*-tolyl-acetamide;
N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(4-fluoro-phenyl)-acetamide;
N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-3,3-dimethyl-butylamide;
N-(2,4-Dimethyl-6-morpholin-4-yl-pyridin-3-yl)-2-(3-fluoro-phenyl)-acetamide;
 2-Bicyclo[2.2.1]hept-2-yl-*N*-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide;
 2-(3,4-Difluoro-phenyl)-*N*-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide;
 4-Methyl-pentanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide;
 2-Cyclopent-2-enyl-*N*-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide;
 2-Cyclohexyl-*N*-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide;
 5-Methyl-hexanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide;
 2-Cyclopentyl-*N*-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-acetamide;
 3-Cyclopentyl-*N*-(2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-propionamide; and
 Hexanoic acid (2,4-dimethyl-6-morpholin-4-yl-pyridin-3-yl)-amide
N-(4-Chloro-2-methoxy-6-morpholin-4-yl-pyridin-3-yl)-2-cyclopentylacetamide
N-(2-Chloro-4-methoxy-6-morpholin-4-yl-pyridin-3-yl)-2-cyclopentylacetamide
N-(2-Chloro-4-methoxy-6-morpholin-4-yl-pyridin-3-yl)-3,3-dimethylbutylamide
N-(4-Chloro-2-methoxy-6-morpholin-4-yl-pyridin-3-yl)-3,3-dimethylbutylamide
N-(4-Chloro-2-methoxy-6-morpholin-4-yl-pyridin-3-yl)-propionamide
 as the free base or salts thereof.

12. A pharmaceutical composition comprising one or more pharmaceutically acceptable carriers or diluents and a compound of the general formula I:



15. Use according to Claim 14 wherein the disorder or disease to be treated is selected from the group consisting of seizure disorders, anxiety disorders, neuropathic pain and migraine pain disorders, neurodegenerative disorders, stroke, cocaine abuse, nicotine withdrawal, ethanol withdrawal and tinnitus.
- 5
16. Use according to Claim 15 wherein the seizure disorders are selected from the group consisting of acute seizures, convulsions, status epilepticus, epilepsy such as epileptic syndromes and epileptic seizures.
- 10
17. Use according to Claim 15 wherein the anxiety disorders are selected from the group consisting of anxiety and disorders and diseases related to panic attack, agoraphobia, panic disorder with agoraphobia, panic disorder without agoraphobia, agoraphobia without history of panic disorder, specific phobia, social phobia and other specific phobias, obsessive-compulsive disorder, post-traumatic stress disorder, acute stress disorders, generalized anxiety disorder, anxiety disorder due to general medical condition, substance-induced anxiety disorder, separation anxiety disorder, adjustment disorders, performance anxiety, hypochondriacal disorders, anxiety disorder due to general medical condition and substance-induced anxiety disorder and anxiety disorder not otherwise specified.
- 15
- 20
18. Use according to Claim 15 wherein the neuropathic pain and migraine pain disorders are selected from the group consisting of allodynia, hyperalgesic pain, phantom pain, neuropathic pain related to diabetic neuropathy, neuropathic pain related to trigeminal neuralgia and neuropathic pain related to migraine.
- 25
19. Use according to Claim 15 wherein the neurodegenerative disorders are selected from the group consisting of Alzheimer's disease, Huntington's chorea, multiple sclerosis, amyotrophic lateral sclerosis, Creutzfeld-Jakob's disease, Parkinson's disease, encephalopathies induced by AIDS or infection by rubella viruses, herpes viruses, borrelia or unknown pathogens, trauma-induced neurodegenerations, neuronal hyperexcitation states such as in medicament withdrawal or intoxication and neurodegenerative diseases of the peripheral nervous system such as polyneuropathies and polyneuritides.
- 30

20. Use according to Claim 14 wherein the disorder or disease to be treated is selected from the group consisting of bipolar disorders.
- 5 21. Use according to Claim 14 wherein the disorder or disease to be treated is selected from the group consisting of sleep disorders; such as insomnia

INTERNATIONAL SEARCH REPORT

International application No
PCT/DK2006/000123

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D413/04 A61K31/5377 A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, EMBASE, BEILSTEIN Data, CHEM ABS Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 02/066036 A (BRISTOL-MYERS SQUIBB COMPANY) 29 August 2002 (2002-08-29) Claims 1-6; Formula (I); examples 1-2, 13, 16, 18, 22, 25 and 30	1-21
Y	WO 01/10380 A (ICAGEN, INC.; MCNAUGHTON-SMITH, GRANT, ANDREW; GROSS, MICHAEL, FRANCIS;) 15 February 2001 (2001-02-15) Claims 1-48; Formula (I); examples 1-9; Fig. 1	1-21
A	WO 2004/082677 A (H. LUNDBECK A/S; KHANZHIN, NIKOLAY; ROTTLAENDER, MARIO; RITZEN, ANDREA) 30 September 2004 (2004-09-30) Claims 1-36; Formula (I); examples	1-21

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

12 June 2006

Date of mailing of the international search report

22/06/2006

Name and mailing address of the ISA/
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kirsch, C

INTERNATIONAL SEARCH REPORT

International application No

PCT/DK2006/000123

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, Y	<p>WO 2005/087754 A (H. LUNDBECK A/S; WENZEL TORNOEE, CHRISTIAN; ROTTLAENDER, MARIO; KHANZH) 22 September 2005 (2005-09-22) Claims 1-20; Formula (I); examples</p>	1-21

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK2006/000123

Box II Observations where certain claims were found unsearchable (Continuation of Item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 13-21 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of Item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/DK2006/000123

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 02066036	A	29-08-2002	CA 2438231 A1	29-08-2002
			CZ 20032233 A3	15-12-2004
			EP 1361879 A1	19-11-2003
			HU 0303841 A2	01-03-2004
			JP 2005508833 T	07-04-2005
			MX PA03007395 A	04-12-2003
			NO 20033683 A	17-10-2003
WO 0110380	A	15-02-2001	AU 779034 B2	06-01-2005
			AU 6758500 A	05-03-2001
			CA 2378241 A1	15-02-2001
			EP 1208085 A2	29-05-2002
			JP 2003506387 T	18-02-2003
			NZ 516610 A	28-05-2004
			ZA 200200502 A	05-02-2003
WO 2004082677	A	30-09-2004	AU 2004222626 A1	30-09-2004
			BR PI0408437 A	04-04-2006
			CA 2519582 A1	30-09-2004
			EP 1613303 A1	11-01-2006
			MX PA05010000 A	17-11-2005
WO 2005087754	A	22-09-2005	NONE	